

Higher covariant derivative Pauli-Villars regularization does not lead to a consistent QCD

C. P. Martin

*Departamento de Física Teórica, C-XI, Universidad Autónoma de Madrid,
Cantoblanco, Madrid 28049, Spain*

F. Ruiz Ruiz

NIKHEF-H, Postbus 41882, 1009 DB Amsterdam, The Netherlands

We compute the beta function at one loop for Yang-Mills theory using as regulator the combination of higher covariant derivatives and Pauli-Villars determinants proposed by Faddeev and Slavnov. This regularization prescription has the appealing feature that it is manifestly gauge invariant and essentially four-dimensional. It happens however that the one-loop coefficient in the beta function that it yields is not $-11/3$, as it should be, but $-23/6$. The difference is due to unphysical logarithmic radiative corrections generated by the Pauli-Villars determinants on which the regularization method is based. This no-go result discards the prescription as a viable gauge invariant regularization, thus solving a long-standing open question in the literature. We also observe that the prescription can be modified so as to not generate unphysical logarithmic corrections, but at the expense of losing manifest gauge invariance.

1 Introduction

To regularize and renormalize a theory with symmetries, it is very convenient to use a regularization method and a renormalization scheme that preserve the symmetries of the theory. In the case of gauge theories, there are not so many regularization methods that preserve gauge invariance. The two traditional candidates are dimensional regularization [1] and higher covariant derivatives [2] [3] –more recent regularization methods [4] [5] will not concern us here. Dimensional regularization works well for vector gauge theories like Yang-Mills or QCD but presents some problems when it comes to dealing with theories with γ^5 . As regards higher covariant derivative regularization, it has the appealing feature that it does not have problems with γ^5 , thus constituting an in principle useful tool to study anomalies. Furthermore, there are suggestions [6] that it leads to supersymmetric as well as gauge invariant regularization procedures. Yet it is not known whether higher covariant derivative regulators work even for the simplest cases, like ordinary Yang-Mills or QCD. In this paper we provide an answer to this question in the negative for the case of Yang-Mills theory.

So let us consider Yang-Mills theory. Higher covariant derivative regularization as proposed in ref. [3] combines a higher covariant derivative term $(D^2 F)^2/4\Lambda^4$ with a set of gauge invariant Pauli-Villars determinants. There is some controversy in the literature as to whether or not this prescription works. On the one side there is the position defended in ref. [3], that claims that the prescription works to all orders in perturbation theory without problems. On the other there is the position of ref. [7], that states that the prescription only regularizes one-loop divergences and leaves two and higher loops unregularized. Even worse, ref. [8] sustains that already at one loop the divergences that arise in the regularized Green functions when the regulators are removed can not be subtracted in a gauge invariant fashion, so that the resulting renormalized theory is not gauge invariant. This conclusion is in turn somewhat in contradiction with the non-multiplicative renormalization schemes discussed in ref. [9]. The problem is that many of these claims, if not all, are based on rather qualitative arguments and are not supported by explicit calculations, thus making it difficult to conclude about the viability of the prescription. To the best of our knowledge, nowhere in the literature physical issues like unitarity or if the prescription reproduces asymptotic freedom with the correct beta function are addressed.

In this paper we use the prescription to explicitly regularize and renormalize Yang-Mills theory at one loop and to compute the beta function at this order in perturbation theory.

It turns out that the renormalized theory is gauge invariant, thus giving lie to ref. [8], but it also turns out that the Pauli-Villars determinants generate unphysical logarithmic radiative corrections. These unphysical corrections modify the beta function of the theory so that the one-loop coefficient in the beta function comes out to be $-23/6$ and not $-11/3$, as it should. In Minkowski space these extra corrections change the imaginary part of some 1PI Green functions. This is in contradiction with general results from renormalization theory according to which the one-loop imaginary part of a 1PI Green function is independent of the renormalization procedure. All this implies that the resulting renormalized theory is inconsistent and that the prescription is not an acceptable one and must be disregarded.

The paper is organized as follows. In section 2 we review the regularization prescription. We spend some time on this since there are a few issues which have passed unnoticed in the previous literature and they play a crucial part in the outcome of this investigation. In section 3 we show that it is possible to subtract the divergences from the regularized theory and have at the same time gauge invariance. In section 4 we compute the beta function at one loop and obtain that it does not have the correct coefficient. Section 5 is devoted to prove that the source of the unwanted contributions to the beta function is the Pauli-Villars determinants on which the prescription is based. We include two appendices which contain the Feynman rules (Appendix A) and calculational details (Appendix B).

2 The regularization method

Let us start introducing our notation and conventions. The theory we are interested in is Yang-Mills theory in four-dimensional Euclidean space. The classical action in a general covariant gauge has the form

$$S = S_{\text{YM}} + S_{\text{GF}} , \quad (2.1)$$

where S_{YM} and S_{GF} are given by

$$S_{\text{YM}} = \frac{1}{4} \int d^4x F_{\mu\nu}^a F^{a\mu\nu} \quad (2.2)$$

and

$$S_{\text{GF}} = \int d^4x \left[\frac{\alpha}{2} b^a b^a - b^a (\partial_\mu A^{a\mu}) + \bar{c}^a (\partial_\mu D^\mu c)^a \right] . \quad (2.3)$$

Here $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c$ denotes the field strength, A_μ^a the gauge field, g the coupling constant and f^{abc} the structure constants of the gauge algebra. The parameter

α is the gauge-fixing parameter, b^a is the auxiliary field, $\bar{c}^a$ and c^a are the Faddeev-Popov ghosts and $D_\mu^{ac} = \delta^{ac} + g f^{abc} A_\mu^b$ is the covariant derivative. The Landau gauge corresponds to $\alpha = 0$, and the Feynman gauge to $\alpha = 1$. We will assume that the gauge group is a compact, simple Lie group so that the structure constants f^{abc} can be taken completely antisymmetric in their indices without loss of generality. Furthermore, we will normalize f^{abc} so that $f^{acd} f^{bcd} = c_V \delta^{ab}$, with c_V the quadratic Casimir operator in the adjoint representation [$c_V = N$ for a gauge group $SU(N)$]. After gauge fixing, gauge invariance takes the form of BRS invariance and so the symmetry to keep under control is the BRS symmetry. The BRS transformations read in our notation

$$sA_\mu^a = (D_\mu c)^a \quad sb^a = 0 \quad s\bar{c}^a = b^a \quad sc^a = -\frac{1}{2} g f^{abc} c^b c^c \quad (2.4)$$

with the BRS operator s satisfying as usual $s^2 = 0$. Finally, the generating functional is formally given by

$$Z[J, \chi, \eta, \bar{\eta}] = \int \mathcal{D}A \mathcal{D}b \mathcal{D}\bar{c} \mathcal{D}c e^{-(S+S_J)},$$

where S is as in eqs. (2.1)-(2.3), S_J is the source term

$$S_J = \int d^4x \left(J^{a\mu} A_\mu^a + \chi^a b^a + \bar{c}^a \eta^a + \bar{\eta}^a c^a \right)$$

and $J^{a\mu}$, χ^a , η^a and $\bar{\eta}^a$ are sources for the fields A_μ^a , b^a , $\bar{c}^a$ and c^a , respectively.

Since it will be used as a point of reference below, let us recall how power counting goes. Given a 1PI Feynman diagram with E_A external gauge lines, E_g external ghost lines and L loops, its overall UV degree ω is given by

$$\omega = 4 - E_A - \frac{3}{2} E_g. \quad (2.5)$$

Superficially divergent 1PI diagrams have $\omega \geq 0$ and correspond to Green functions already present in the classical action S . As is well known, this ensures renormalizability. To actually renormalize the theory, one proceeds in two steps. The first one is to provide a regularization method that renders all Green functions finite order by order in perturbation theory. The second one is to provide a renormalization scheme that eliminates the divergences appearing in the Green functions when the regulator is removed. In this paper we use higher covariant derivative Pauli-Villars regularization as proposed by Faddeev and Slavnov [2] [3] to explicitly renormalize Yang-Mills theory at one loop. As announced in the introduction, it will turn out that the resulting renormalized theory is inconsistent. In the course of our investigation we have found that there are a few issues concerning the regularization method itself that

have gone unnoticed in the previous literature and that play a crucial role in the result. We thus find worth spending some time in reviewing the method and bringing these points to light.

2.1 Higher covariant derivative regulators

The method proposed by Faddeev and Slavnov regularizes Yang-Mills theory in two steps. In the first one, it introduces a higher covariant derivative mass Λ by replacing the classical gauge-fixed action S in eq. (2.1) with the action [3]

$$S_\Lambda = S_{\text{YM}} + S_2 + S_{\text{GF}}^f, \quad (2.6)$$

where S_{YM} is as in eq. (2.2) and S_2 and S_{GF}^f are respectively given by

$$S_2 = \frac{1}{4\Lambda^4} \int d^4x (D^2 F_{\mu\nu})^a (D^2 F^{\mu\nu})^a \quad (2.7)$$

and

$$S_{\text{GF}}^f = \int d^4x \left[\frac{\alpha}{2} b^a \frac{1}{f^2(\partial^2/\Lambda^2)} b^a - b^a (\partial_\mu A^{a\mu}) + \bar{c}^a (\partial_\mu D^\mu c)^a \right].$$

Here $f(\partial^2/\Lambda^2)$ is a polynomial in its argument that in momentum space becomes a function of p^2/Λ^2 and that by definition is chosen to satisfy the conditions

$$\begin{aligned} f(p^2/\Lambda^2) &\rightarrow 1 \quad \text{as } \Lambda \rightarrow \infty \\ f(p^2/\Lambda^2) &\sim (p^2/\Lambda^2)^\gamma \quad \text{as } |p| \rightarrow \infty, \end{aligned} \quad (2.8)$$

with $\gamma \geq 1$. In this paper we will take

$$f\left(\frac{p^2}{\Lambda^2}\right) = 1 + \frac{p^4}{\Lambda^4} \quad (2.9)$$

for reasons that we explain below. The propagator of the gauge field for the action S_Λ has the form

$$D_{\mu\nu}^{ab}(p) = \delta^{ab} \left[\frac{\Lambda^4}{p^4(p^4 + \Lambda^4)} (p^2 g_{\mu\nu} - p_\mu p_\nu) + \alpha \frac{p_\mu p_\nu}{p^4 f^2(p^2/\Lambda^2)} \right], \quad (2.10)$$

where p^{2n} is a shorthand notation for $(p^2)^n$. The second condition in eq. (2.8) ensures that the propagator goes as $1/p^6$ when $p^\mu \rightarrow \infty$. This and some power counting implies that the overall UV degree ω_Λ of a 1PI diagram generated by S_Λ with E_A external gauge legs, E_g external ghost legs and L loops is given by [compare with eq. (2.5)]

$$\omega_\Lambda = 4 - 4(L - 1) - E_A - \frac{7}{2} E_g. \quad (2.11)$$

Now only one-loop 1PI diagrams with $E_A = 2, 3, 4$ and $E_g = 0$ are superficially divergent. We thus see that the higher covariant derivative regulator Λ improves the power counting behaviour of the theory but does not completely regularize it. To achieve a complete regularization, one must regularize the one-loop diagrams for which $\omega_\Lambda \geq 0$. Before doing so, let us make a few comments that we feel are in order.

Comment 1. There are many possible choices for f compatible with conditions (2.8). The choice in eq. (2.9) ensures that α -dependent contributions in any 1PI diagram are finite by power counting, as can be trivially checked. In fact, any f for which γ in eq. (2.8) is larger or equal than 2 leads to superficially convergent α -dependent contributions. We have taken f as in eq. (2.9) since in this way the denominator of the propagator $D_{\mu\nu}^{ab}(p)$ has factors only of type p^4 and $p^4 + \Lambda^4$. Note also that we have required f to be polynomial so as to ensure locality.

Comment 2. Since the action S_Λ is BRS invariant, the higher covariant derivative regulator Λ preserves BRS invariance.

Comment 3. Note that all 1PI diagrams generated by S_Λ with external ghost legs are finite by power counting. This implies that Yang-Mills 1PI diagrams with external ghosts are fully regularized by the higher covariant derivative regulator Λ .

Comment 4. One may wonder whether the theory can be completely regularized by means of a higher covariant derivative term of higher order in the covariant derivative. The answer is no. To see this, consider the term

$$S_n = \frac{(-)^n}{4\Lambda^{2n}} \int d^4x F_{\mu\nu}^a (D^{2n} F^{\mu\nu})^a,$$

where n is an arbitrary positive integer. Power counting now implies that

$$\omega_\Lambda(n) = 4 - 2n(L - 1) - E_A - \left(\frac{3}{2} + n\right)g.$$

So no matter how large n is, $\omega_\Lambda(n)$ is always negative for $L = 1$, $E_g = 0$ and $E_A = 2, 3, 4$.

2.2 Pauli-Villars determinants

To regularize the one-loop divergences in the Feynman diagram expansion generated by the action S_Λ above, one can proceed as follows [2] [3]. Consider the two following generating

functionals labeled by an index $r = 0, 1$:

$$Z_r[J, \chi, \eta, \bar{\eta}] = \int \mathcal{D}A \mathcal{D}b \mathcal{D}\bar{c} \mathcal{D}c e^{-(S_\Lambda + S_J)} \prod_{j=1}^J (\det \mathbb{A}_j^r)^{-\alpha_j/2} \prod_{i=1}^I (\det \mathbb{C}_i^r)^{\gamma_i}, \quad (2.12)$$

Here $\{\alpha_j\}_J$ and $\{\gamma_i\}_I$ are real parameters to be fixed and $\det \mathbb{A}_j^r$ and $\det \mathbb{C}_i^r$ are defined by

$$(\det \mathbb{A}_j^r)^{-1/2} = \int \mathcal{D}A_j \mathcal{D}b_j \exp \left\{ - (S_{M_j} + S_{b_j}) \right\} \quad (2.13)$$

and

$$\det \mathbb{C}_i^r = \int \mathcal{D}\bar{c}_j \mathcal{D}c_j \exp \left\{ - S_{m_i} \right\}, \quad (2.14)$$

with

$$\begin{aligned} S_{M_j} &= \frac{1}{2} \int d^4x \int d^4y A_{j\mu}^a(x) \left[\frac{\delta^2 S_\Lambda}{\delta A_\mu^a(x) \delta A_\nu^b(y)} - M_j^2 \delta^{ab} g^{\mu\nu} \delta(x-y) \right] A_{j\nu}^b(y) \\ S_{b_j} &= - \int d^4x b_j^a (\delta^{ab} \partial_\mu + r g f^{acb} A_\mu^c) A_j^{b\mu} \\ S_{m_i} &= \int d^4x \left[\bar{c}_i^a (\partial_\mu D^\mu c_i)^a + r g f^{abc} \bar{c}_i^a A_\mu^b (D^\mu c_i)^c - m_i^2 \bar{c}_i^a c_i^a \right]. \end{aligned}$$

The fields $\{A_{j\mu}^a\}$ are commuting Pauli-Villars fields of mass $\{M_j\}$, the fields b_j^a are Pauli-Villars Lagrange multipliers imposing the condition

$$(\delta^{ab} \partial_\mu + r g f^{acb} A_\mu^c) A_j^{b\mu} = 0$$

and the fields $\{\bar{c}_i^a, c_i^a\}$ are anticommuting Pauli-Villars fields of mass $\{m_i\}$. The determinants $\det \mathbb{A}_j^r$ and $\det \mathbb{C}_i^r$ are BRS invariant if $r = 1$ –actually gauge invariant since they do not depend on ghosts. This follows from the observation [3] that the change of integration variables $\phi \rightarrow \phi + \theta(s\phi)$, where θ is an anticommuting parameter and $s\phi$ is given by

$$sA_{j\mu}^a = g f^{abc} A_{j\mu}^b c^c \quad sb_j^a = g f^{abc} b_j^b c^c \quad s\bar{c}_i^a = g f^{abc} \bar{c}_i^b c^c \quad sc_i^a = g f^{abc} c_i^b c^c,$$

cancels the extra terms that arise in $\det \mathbb{A}_j^1$ and $\det \mathbb{C}_i^1$ when the gauge field A_μ^a is BRS-transformed. However, if $r = 0$, the determinants $\det \mathbb{A}_j^r$ and $\det \mathbb{C}_i^r$ are not BRS invariant. Faddeev's and Slavnov's proposal [2] [3] corresponds to taking $r = 1$ and is the one we will be studying in this paper. For later convenience, though, we will keep r explicit in our calculations.

The Feynman rules associated to the functional $Z_r[J, \chi, \eta, \bar{\eta}]$ are listed in Appendix A. Note that the determinant $\det \mathbb{A}_j^r$ has been defined in such a way that the vertices

A^{2l+1} ($l = 3, \dots, 6$) have the same Feynman rule as $A_j^2 A^l$. Note also that the Feynman rules for the vertices $b_j A A_j$, $\bar{c}_i A c_i$ and $\bar{c}_i A^2 c_i$ depend explicitly on r . Keeping these two observations in mind, we move on to discussing how the Pauli-Villars determinants just introduced regularize the one-loop divergences generated by S_Λ . We start discussing first the case $r = 0$.

Gauge non-invariant case: $r = 0$

We want to show that the vacuum polarization tensor and the vertices $\langle A^3 \rangle_{\text{1PI}}$ and $\langle A^4 \rangle_{\text{1PI}}$ at one loop generated by $Z_0[J, \chi, \eta, \bar{\eta}]$ in eq. (2.12) are finite. Let us start with the vacuum polarization tensor. At one loop it receives contributions from the diagrams in Fig. 1. Let us look at the sum of the diagrams in Figs. (1a) and (1b),

$$\sigma_1 = (1a) + (1b). \quad (2.15)$$

Recalling that α -dependent contributions are superficially convergent, we have that in diagram (1a) only the $\alpha = 0$ part contains divergences. Taking then $\alpha = 0$, noting that by construction the vertices A^3 and $A^2 A_j^2$ have the same Feynman rule, and using the identity

$$\frac{1}{q^6 + q^2 \Lambda^4 + M_j^2 \Lambda^4} = \frac{1}{q^2 (q^4 + \Lambda^4)} - \frac{M_j^2 \Lambda^4}{q^2 (q^4 + \Lambda^4) (q^6 + q^2 \Lambda^4 + M_j^2 \Lambda^4)} \quad (2.16)$$

for the propagator of the Pauli-Villars field A_j , it is trivial to see that the only superficially divergent integrals in σ_1 are quadratically divergent, do not depend on M_j and pick an overall factor $1 + \sum_j \alpha_j$. It thus follows that if we choose parameters α_j such that

$$1 + \sum_{j=1}^J \alpha_j = 0 \quad (2.17)$$

the sum σ_1 becomes finite. In other words, diagram (1b) regularizes diagram (1a) if condition (2.17) is satisfied. A similar argument shows that the sum

$$\sigma_2 = (1c) + (1d) \quad (2.18)$$

is finite if again condition (2.17) is met. Let us finally consider the sum

$$\sigma_3^0 = (1e) + (1f). \quad (2.19)$$

Noting that for $r = 0$ the vertex $\bar{c}_i A c_i$ has the same Feynman rule as the vertex $\bar{c} A c$ and using the identity

$$\frac{1}{q^2 + m_i^2} = \frac{1}{q^2} - \frac{m_i^2}{q^2 (q^2 + m_i^2)}$$

twice, we find that σ_3^0 contains quadratic and logarithmic divergences that do not depend on m_i and that pick overall factors $1 + \sum_i \gamma_i$ and $\sum_i \gamma_i m_i^2$, respectively. Hence imposing conditions

$$1 + \sum_{i=1}^I \gamma_i = 0 \quad (2.20)$$

and

$$\sum_{i=1}^I \gamma_i m_i^2 = 0, \quad (2.21)$$

we render the sum (e) + (f) finite.

To summarize, if the parameters α_j and γ_i and the masses M_j and m_i satisfy the Pauli-Villars conditions (2.17)-(2.21), diagrams (1b), (1d) and (1f) regularize diagrams (1a), (1c) and (1e) and the vacuum polarization tensor becomes finite. Analogous arguments show that one-loop divergences in the vertices $\langle A^3 \rangle_{\text{1PI}}$ and $\langle A^4 \rangle_{\text{1PI}}$ generated by S_Λ are regularized if the very same conditions (2.17)-(2.21) hold. This regularization procedure is very neat, but as already pointed out is not gauge invariant.

Gauge invariant case: $r = 1$

We know that the Feynman rules are not the same for $r = 1$ as for $r = 0$. Hence there is no reason to assume that the regularization mechanism is the same in both cases. To study regularization for $r = 1$ in detail, let us consider again the vacuum polarization tensor. At one loop, it receives contributions from the diagrams in Fig. 1 and, very importantly, from those in Fig. 2. We first look at the diagrams in Fig. 1. Diagrams (1a) to (1d) are independent of r , so the same arguments as for $r = 0$ imply that the sums σ_1 and σ_2 are finite provided condition (2.17) holds. As concerns the sum σ_3^0 , we note that diagram (1e) is independent of r but (1f) is not. Furthermore, the difference in between taking $r = 1$ and $r = 0$ in diagram (1f) is a superficially divergent integral. Therefore, after imposing conditions (2.20) and (2.21), we are left with unregularized divergences in σ_3^0 . Next we look at the diagrams in Fig. 2, which we emphasize do not exist for $r = 0$. Some trivial power counting shows that diagram (2a) is finite, while diagrams (2b) to (2e) are superficially divergent. Hence, for the generating functional $Z_1[J, \chi, \eta, \bar{\eta}]$ to produce a finite vacuum polarization tensor, the divergences in (1e) and (1f) must cancel the divergences in diagrams (2b) to (2e). In other words, the sum

$$\sigma_3^1 = (1e) + (1f) + (2b) + (2c) + (2d) + (2e) \quad (2.22)$$

must be finite. All this makes obvious that regularization for $r = 1$ does not go as for $r = 0$. It also corrects some statements in the previous literature [3], where it is explicitly claimed that the terms that ensure manifest BRS invariance do not introduce new divergences into the game.

Let us find the conditions on α_j , γ_i , M_j and m_i for which the sum σ_3^1 is finite. A look at the topology of the diagrams contributing to σ_3^1 reveals that an analysis of the type performed for σ_1 and σ_2 is not possible. Yet we have to seek for cancellations of divergences from different diagrams. To do this with rigour, we introduce a regulator $\mathcal{R}$, compute the diagrams at finite $\mathcal{R}$ and see for what values of α_j , γ_i , M_j and m_i contributions divergent at $\mathcal{R} = 0$ cancel upon summation. To isolate the divergences at $\mathcal{R} = 0$ in each one of the diagrams, we proceed as follows. Consider for example diagrams (2c) and (2d). With the choice of momenta in Fig. 2 and after some Lorentz algebra¹, we have

$$(2c) + (2d) = r g^2 c_V \sum_{j=1}^J \alpha_j \int_{\mathcal{R}} \frac{d^4 q}{(2\pi)^4} \frac{N_{\mu\nu}}{(q+p)^2 q^2 (q^6 + q^2 \Lambda^4 + M_j^2 \Lambda^4)} ,$$

where

$$N_{\mu\nu} = 2 q^6 (q_\mu q_\nu - q^2 g_{\mu\nu}) + O(q^n, 2 \leq n < 6) . \quad (2.23)$$

Terms in $N_{\mu\nu}$ of order five or less in q^μ give rise to integrals finite by power counting at $\mathcal{R} = 0$. On the other hand, terms in $N_{\mu\nu}$ of order six or higher in q^μ give rise to superficially divergent integrals at $\mathcal{R} = 0$. This means that as far as regularization is concerned it is enough to keep the terms written explicitly in eq. (2.23). Using then the identities (2.16) and

$$\frac{1}{q^4 + \Lambda^4} = \frac{1}{q^4} - \frac{1}{q^4(q^4 + \Lambda^4)} \quad (2.24)$$

and retaining only integrals which are superficially divergent at $\mathcal{R} = 0$, it is trivial to see that

$$(2c) + (2d) \Big|_{\text{div}} = -r g^2 c_V \delta^{ab} \sum_{j=1}^J \alpha_j \int_{\mathcal{R}} \frac{d^4 q}{(2\pi)^4} \frac{2 (g_{\mu\nu} - q_\mu q_\nu)}{(p+q)^2} .$$

Proceeding in the same way for the other diagrams in σ_3^1 , summing over diagrams and

¹To perform the Lorentz algebra of the diagrams in this paper, we have used the algebraic package REDUCE [10]

imposing condition (2.17), we obtain

$$\begin{aligned}
\sigma_3^1(\mathcal{R}) \Big|_{\text{div}} &= -g^2 c_V \delta^{ab} \left(1 + \sum_{i=1}^I \gamma_i \right) \int_{\mathcal{R}} \frac{d^4 q}{(2\pi)^4} \frac{4 q_\mu q_\nu + q_\mu p_\nu + p_\mu q_\nu}{q^2 (q+p)^2} \\
&\quad + 2 g^2 c_V \delta^{ab} \int_{\mathcal{R}} \frac{d^4 q}{(2\pi)^4} \left\{ \sum_{i=1}^I \gamma_i \left[\frac{g_{\mu\nu}}{q^2 + m_i^2} + \frac{4 m_i^2 q_\mu q_\nu}{q^4 (q^2 + m_i^2)} \right] + \frac{g_{\mu\nu}}{(q+p)^2} \right\} \\
&\quad - g^2 c_V \delta^{ab} \sum_{i=1}^I \gamma_i \int_{\mathcal{R}} \frac{d^4 q}{(2\pi)^4} \frac{q_\mu p_\nu + p_\mu q_\nu + p_\mu p_\nu}{q^2 (q+p)^2}.
\end{aligned} \tag{2.25}$$

Assume we take as regulator $\mathcal{R}$ dimensional regularization, *i.e.* assume that

$$\int_{\mathcal{R}} \frac{d^4 q}{(2\pi)^4} = \frac{1}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \quad D = 4 + 2\varepsilon,$$

with μ the dimensional regularization mass scale. Then, it is not difficult to see that

$$\sigma_3^1(\varepsilon) \Big|_{\text{div}} = \frac{g^2 c_V}{16\pi^2} \delta^{ab} \left[\left(1 + \sum_{i=1}^I \gamma_i \right) \sigma_\varepsilon (p^2 g_{\mu\nu} - p_\mu p_\nu) + \sum_{i=1}^I \gamma_i p^2 g_{\mu\nu} \right] \tag{2.26}$$

as $D \rightarrow 4$, where

$$\sigma_\varepsilon = \frac{1}{3} \left[\frac{1}{\varepsilon} + \ln\left(\frac{p^2}{\mu^2}\right) + \gamma - \frac{8}{3} \right].$$

To get rid of the pole in eq. (2.26) and end up with a finite σ_3^1 , we choose the parameters γ_i to satisfy the Pauli-Villars condition (2.20).

Putting everything together, we have that the regularized vacuum polarization tensor is finite for parameters α_j and γ_i satisfying the Pauli-Villars conditions (2.17) and (2.20). Using the BRS identities and that the vacuum polarization tensor, the ghost self-energy and the $\bar{c}Ac$ -vertex are finite, it follows the three and four-point Green functions $\langle A^3 \rangle_{\text{1PI}}$ and $\langle A^4 \rangle_{\text{1PI}}$ are also finite, provided of course conditions (2.17) and (2.20) hold. This closes the proof that the functional $Z_1[J, \chi, \eta, \bar{\eta}]$ generates finite one-loop Green functions. There are two important comments to be made at this point:

Comment 1. It is clear that regularization does not require condition (2.21), contrarily to what is claimed in ref. [3] and all papers thereafter.

Comment 2. Divergences from isolated Feynman diagrams in σ_3^1 get regularized only after integration at finite $\mathcal{R}$. This is in contrast with what happens for σ_1 and σ_2 , where regularization takes place algebraically prior to integration. Here we have taken dimensional

regularization as regulator $\mathcal{R}$ because it preserves BRS invariance. Assume that instead we take a cut-off Q such that $|p|, m_i \ll Q$. Some elementary integration in eq. (2.25) then leads to

$$\sigma_3^1(Q)|_{\text{div}} = \frac{g^2 c_V}{16\pi^2} \delta^{ab} \left[\left(1 + \sum_{i=1}^I \gamma_i \right) (Q^2 g_{\mu\nu} + \sigma'_Q p^2 g_{\mu\nu} - \sigma''_Q p_\mu p_\nu) + \sum_{i=1}^I \gamma_i \left(p^2 g_{\mu\nu} - \frac{1}{2} p_\mu p_\nu \right) \right] \quad (2.27)$$

where

$$\sigma'_Q = \frac{1}{3} \ln\left(\frac{Q^2}{p^2}\right) + \frac{11}{18} \quad \sigma''_Q = \frac{1}{3} \ln\left(\frac{Q^2}{p^2}\right) - \frac{1}{18}.$$

Again we see that for γ_i satisfying eq. (2.20), the sum σ_3^1 is finite. Comparing eq. (2.26) with eq. (2.27), we note that the finite part that is left in $\sigma_3^1|_{\text{div}}$ after imposing condition (2.20) is however different. This implies that different $\mathcal{R}$'s give different local contributions to σ_3^1 . As regards the non-local part of σ_3^1 , it is the same for all $\mathcal{R}$ since it arises only from integrals which are finite by power counting at $\mathcal{R} = 0$.

To sum up, Pauli-Villars determinants with $r = 1$ are BRS invariant, but checking that they provide the necessary regularization requires introducing a pre-regulator $\mathcal{R}$. This means strictly speaking that $r = 1$ Pauli-Villars determinants do not provide an acceptable regularization. Still one could argue that since non-local contributions are independent of the choice of pre-regulator $\mathcal{R}$, $r = 1$ Pauli-Villars regularization is good enough. In sections 4 and 5 we show that this is not either the case, since the determinants $\det \mathbb{A}_j^1$ and $\det \mathbb{C}_i^1$ introduce unphysical logarithmic radiative corrections.

3 The renormalized theory

So far we have a BRS invariant regularization prescription that involves the masses Λ , M_j and m_i (and a pre-regulator $\mathcal{R}$ which from now on we take it to be dimensional regularization). The next step is to define the renormalized theory. There are several requirements that the resulting renormalized theory must satisfy for it to make sense. One of them is that the renormalized Green functions satisfy the BRS identities. To date, there is no agreement in the literature [8] as whether starting from the generating functional $Z_1[J, \chi, \eta, \bar{\eta}]$ it is possible to define renormalized Green functions compatible with BRS invariance. In this section we show that this is indeed possible at one loop in the least.

To study the BRS identities, it is convenient to work with the effective action. The effective action associated to the functional $Z_1[J, \chi, \eta, \bar{\eta}]$ is defined in two steps. One first

introduces external fields $K^{a\mu}$ and H^a coupled respectively to the non-linear BRS transforms sA_μ^a and sc^a so that the generating functional becomes

$$Z_1[J, \chi, \eta, \bar{\eta}; K, H] = \int \mathcal{D}A \mathcal{D}b \mathcal{D}\bar{c} \mathcal{D}c e^{-(S_\Lambda + S_J + S_{\text{ext}})} \prod_{j=1}^J (\det \mathbb{A}_j^1)^{-\alpha_j/2} \prod_{i=1}^I (\det \mathbb{C}_i^1)^{\gamma_i},$$

with S_{ext} given by

$$S_{\text{ext}} = \int d^4x \left[K^{a\mu}(sA_\mu^a) + H^a(sc^a) \right]. \quad (3.1)$$

And secondly, one writes

$$Z_1[J, \chi, \eta, \bar{\eta}; K, H] = \exp \left(-W[J, \chi, \eta, \bar{\eta}; K, H] \right)$$

and performs a Legendre transformation on $W[J, \chi, \eta, \bar{\eta}; K, H]$ and on the sources $J^{a\mu}$, χ^a , η^a and $\bar{\eta}^a$. The Legendre transform of $W[J, \chi, \eta, \bar{\eta}; K, H]$ is the regularized effective action, a functional of mass dimension four and ghost number zero that depends on the Legendre fields $A_\mu^a, b^a, \bar{c}^a, c^a$ and on the external fields $K^{a\mu}, H^a$ and that generates 1PI Green functions. In what follows we will denote by the letter ψ the set of fields $\{\psi\} = \{A_\mu^a, b^a, \bar{c}^a, c^a, K^{a\mu}, H^a\}$ and write $\Gamma_{\Lambda M_j m_i}[\psi]$ for the regularized effective action, our conventions for the mass dimensions d_ψ and the ghost numbers gh_ψ of the fields being $\{d_\psi\} = \{1, 2, 1, 1, 2, 2\}$ and $\{gh_\psi\} = \{0, 0, -1, 1, -1, -2\}$. It is very easy to see that all 1PI Feynman diagrams with external fields $K^{a\mu}, H^a$ are superficially convergent. This ensures that the manipulations performed above to arrive at the effective action $\Gamma_{\Lambda M_j m_i}[\psi]$ do not introduce divergences.

We will write

$$\Gamma_{\Lambda M_j m_i}[\psi] = \Gamma_\Lambda^{(0)}[\psi] + \Gamma_{\Lambda M_j m_i}^{(1)}[\psi] + O(\hbar^2),$$

where by definition the tree-level contribution $\Gamma_\Lambda^{(0)}[\psi]$ is given by

$$\Gamma_\Lambda^{(0)}[\psi] = S_\Lambda + S_{\text{ext}}$$

and $\Gamma_{\Lambda M_j m_i}^{(1)}[\psi]$ collects all one-loop corrections. Since the action S_Λ and the determinants $\det \mathbb{A}_j^1$ and $\det \mathbb{C}_i^1$ are BRS invariant, $\Gamma_{\Lambda M_j m_i}[\psi]$ satisfies the BRS identities. At tree level and one loop, they take the form

$$\Delta_\Lambda \Gamma_\Lambda^{(0)}[\psi] = 0 \quad (3.2)$$

$$\Delta_\Lambda \Gamma_{\Lambda M_j m_i}^{(1)}[\psi] = 0, \quad (3.3)$$

where Δ_Λ is the Slavnov-Taylor operator for the action $\Gamma_\Lambda^{(0)}[\psi]$. The operator Δ_Λ can be written as

$$\Delta_\Lambda = \Delta + \int d^4x \frac{\delta S_2}{\delta A_\mu^a} \frac{\delta}{\delta K^{a\mu}},$$

where

$$\Delta = \int d^4x \left[\frac{\delta \Gamma^{(0)}}{\delta A_\mu^a} \frac{\delta}{\delta K^{a\mu}} + \frac{\delta \Gamma^{(0)}}{\delta K^{a\mu}} \frac{\delta}{\delta A_\mu^a} + \frac{\delta \Gamma^{(0)}}{\delta c^a} \frac{\delta}{\delta H^a} + \frac{\delta \Gamma^{(0)}}{\delta H^a} \frac{\delta}{\delta c^a} + b^a \frac{\delta \Gamma^{(0)}}{\delta \bar{c}^a} \right] \quad (3.4)$$

is the Slavnov-Taylor operator for Yang-Mills theory and S_2 is as in eq. (2.7), the action $\Gamma^{(0)}[\psi]$ being the tree-level Yang-Mills action

$$\Gamma^{(0)}[\psi] = S_{\text{YM}} + S_{\text{GF}} + S_{\text{ext}}. \quad (3.5)$$

For later use, we recall that Δ is nilpotent, $\Delta^2 = 0$, and satisfies $\Delta S_{\text{YM}} = 0$. Eq. (3.2) is trivially satisfied does not provide any new information. We want to study the structure of the 1PI Green functions that the regularized effective action $\Gamma_{\Lambda M_j m_i}[\psi]$ generates and the constraints that eq. (3.3) imposes on them.

Let us then consider the regularized vacuum polarization tensor $\Pi_{\mu\nu}^{ab}(p, \Lambda, M_j, m_i)$. Manifest BRS invariance implies that up to one loop it will be of the form

$$\Pi_{\mu\nu}^{ab}(p, \Lambda, M_j, m_i) = \left[1 + \frac{g^2 c_V}{16\pi^2} \Pi(p, \Lambda, M_j, m_i) \right] \delta^{ab} (p^2 g_{\mu\nu} - p_\mu p_\nu), \quad (3.6)$$

where $\Pi(p, \Lambda, M_j, m_i)$ is a dimensionless function of its arguments. Since $\Pi(p, \Lambda, M_j, m_i)$ is dimensionless, it can be written without loss of generality as a function of p^2/Λ^2 , p^2/M_j^2 and p^2/m_i^2 . Power counting implies that in the limit $\Lambda, M_j, m_i \rightarrow \infty$, $\Pi(p, \Lambda, M_j, m_i)$ will take the form

$$\Pi(p, \Lambda, M_j, m_i) = A_3 \ln\left(\frac{p^2}{\Lambda^2}\right) + B_3 \sum_{j=1}^J \alpha_j \ln\left(\frac{p^2}{M_j^2}\right) + C_3 \sum_{i=1}^I \gamma_i \ln\left(\frac{p^2}{m_i^2}\right) + \pi_0, \quad (3.7)$$

with A_3 , B_3 , C_3 and π_0 coefficients that depend on α . Assume more generally now that $G^{(0)}(p_e)$ denotes an arbitrary 1PI Yang-Mills Green function at tree level with $\{p_e\} = \{p_1^\mu, \dots, p_E^\mu\}$ independent external momenta. Manifest BRS invariance and power counting imply that in the limit $\Lambda, M_j, m_i \rightarrow \infty$ the corresponding regularized Green function, $G(p_e, \Lambda, M_j, m_i)$, can be written up to one loop without loss of generality as

$$G(p_e, \Lambda, M_j, m_i) = G^{(0)}(p_e) + \frac{g^2 c_V}{16\pi^2} \left[\phi_G(p, \Lambda, M_j, m_i) G^{(0)}(p_e) + G^{\text{fin}}(p_e) \right].$$

where $\phi_G(p, \Lambda, M_j, m_i)$ is given by

$$\phi_G(p, \Lambda, M_j, m_i) = A_G \ln\left(\frac{p^2}{\Lambda^2}\right) + B_G \sum_{j=1}^J \alpha_j \ln\left(\frac{p^2}{M_j^2}\right) + C_G \sum_{i=1}^I \gamma_i \ln\left(\frac{p^2}{m_i^2}\right) \quad (3.8)$$

and $G^{\text{fin}}(p_e)$ collects all finite contributions. The momentum p^μ in eq. (3.8) is a non-zero linear combination of the external momenta and can be chosen arbitrarily since

$$\ln(p^2/\Lambda^2) = \ln(p'^2/\Lambda^2) + \ln(p^2/p'^2)$$

for all $p'^2 \neq 0$. Obviously, different choices of p^μ give different finite parts $G_{\text{fin}}(p_e)$ but the same coefficients A_G , B_G and C_G . Introducing a renormalization mass scale κ and using the Pauli-Villars conditions (2.17) and (2.20), we can cast $G(p_e, \Lambda, M_j, m_i)$ as

$$G(p_e, \Lambda, M_j, m_i) = G^{(0)}(p_e) + \frac{g^2 c_V}{16\pi^2} \left[\phi_G(\kappa, \Lambda, M_j, m_i) G^{(0)}(p_e) + G_\kappa^{\text{fin}}(p_e) \right], \quad (3.9)$$

where now

$$\phi_G(\kappa, \Lambda, M_j, m_i) = A_G \ln\left(\frac{\kappa^2}{\Lambda^2}\right) + B_G \sum_{j=1}^J \alpha_j \ln\left(\frac{\kappa^2}{M_j^2}\right) + C_G \sum_{i=1}^I \gamma_i \ln\left(\frac{\kappa^2}{m_i^2}\right) \quad (3.10)$$

and

$$G_\kappa^{\text{fin}}(p_e) = (A_G - B_G - C_G) \ln\left(\frac{p^2}{\kappa^2}\right) G^{(0)}(p_e) + G^{\text{fin}}(p_e). \quad (3.11)$$

In the sequel we will be using the following notation. Subindices $G = 2, 3, 4$ will respectively refer to the two, three and four-point Green functions $\langle A^2 \rangle_{\text{1PI}}$, $\langle A^3 \rangle_{\text{1PI}}$ and $\langle A^4 \rangle_{\text{1PI}}$; subindices $G = 2$ and a tilde and $G = 3$ and a tilde will refer to $\langle \bar{c}c \rangle_{\text{1PI}}$ and $\langle \bar{c}Ac \rangle_{\text{1PI}}$; and subindices $G = K$ and $G = H$ will refer to $\langle KAc \rangle_{\text{1PI}}$ and $\langle Hcc \rangle_{\text{1PI}}$. Note that since the Pauli-Villars fields do not couple to the Faddeev-Popov ghosts nor to the external fields, we have

$$\tilde{B}_2 = \tilde{B}_3 = 0 \quad \tilde{C}_2 = \tilde{C}_3 = 0 \quad B_K = C_K = 0 \quad B_H = C_H = 0. \quad (3.12)$$

To find the constraints on the coefficients A_G , B_G and C_G that the BRS identity (3.3) imposes, we note that one-loop 1PI diagrams with $K^{a\mu}$ external legs have $B_K = C_K = 0$. Hence these diagrams give contributions to $\Gamma_{\Lambda M_j m_i}^{(1)}[\psi]$ that when $\Lambda, M_j, m_i \rightarrow \infty$ are either finite or diverge as $\ln(\kappa^2/\Lambda^2)$. Recalling that S_2 has a overall factor $1/\Lambda^4$, this implies that

$$\lim_{\Lambda, M_j, m_i \rightarrow \infty} \int d^4x \frac{\delta S_2}{\delta A_\mu^a} \frac{\delta}{\delta K^{a\mu}} \Gamma_{\Lambda M_j m_i}^{(1)}[\psi] = 0. \quad (3.13)$$

From our considerations above it follows that in the limit $\Lambda, M_j, m_i \rightarrow \infty$ the one-loop contribution to the regularized effective action takes the form

$$\Gamma_{\Lambda M_j m_i}^{(1)}[\psi] = \Gamma_{\Lambda M_j m_i \kappa}^{(1) \text{div}}[\psi] + \Gamma_{\kappa}^{(1) \text{fin}}[\psi] + \text{v.t.} , \quad (3.14)$$

where $\Gamma_{\Lambda M_j m_i \kappa}^{(1) \text{div}}[\psi]$ generates the ϕ_G -contributions in eq. (3.10), $\Gamma_{\kappa}^{(1) \text{fin}}[\psi]$ generates finite one-loop radiative corrections and ‘v.t.’ collects contributions that vanish as $\Lambda, M_j, m_i \rightarrow \infty$. Eqs. (3.13) and (3.14) imply that the BRS identity (3.3) reduces in the limit $\Lambda, M_j, m_i \rightarrow \infty$ to

$$\Delta \Gamma_{\Lambda M_j m_i \kappa}^{(1) \text{div}}[\psi] = 0 \quad (3.15)$$

$$\Delta \Gamma_{\kappa}^{(1) \text{fin}}[\psi] = 0 . \quad (3.16)$$

Using that by definition $\Gamma_{\Lambda M_j m_i \kappa}^{(1) \text{div}}[\psi]$ generates the ϕ_G -contributions to the Green functions in eq. (3.10), it is not difficult to see after some algebra that eq. (3.15) implies that the coefficients A_G , B_G and C_G satisfy

$$A_2 - A_3 = A_3 - A_4 = \tilde{A}_2 - \tilde{A}_3 \quad (3.17)$$

$$B_2 = B_3 = B_4 \quad (3.18)$$

$$C_2 = C_3 = C_4 . \quad (3.19)$$

Once we know the form of the divergences in the regularized Green functions when the regulators are sent to infinity, we are in a position to renormalize the theory.

To define a one-loop renormalized Yang-Mills theory consistent with gauge invariance, we have to provide a one-loop renormalized effective action

$$\Gamma_R[\psi] = \Gamma^{(0)}[\psi] + \Gamma_R^{(1)}[\psi] + O(\hbar^2) \quad (3.20)$$

that generates finite 1PI Green functions and that satisfies the Yang-Mills BRS identities. We recall that the latter take the form up to one loop

$$\Delta \Gamma^{(0)}[\psi] = 0 \quad (3.21)$$

$$\Delta \Gamma_R^{(1)}[\psi] = 0 . \quad (3.22)$$

For $\Gamma_R^{(1)}[\psi]$ we take

$$\Gamma_R^{(1)}[\psi] = \Gamma_{\kappa}^{(1) \text{fin}}[\psi] + c_1 S_{\text{YM}} + c_2 \Delta X[\psi] , \quad (3.23)$$

where $\Gamma_{\kappa}^{(1)\text{fin}}[\psi]$ is the same as in eq. (3.14), c_1 and c_2 are arbitrary coefficients of order $\hbar$ and $X[\psi]$ is the most general local integrated functional of mass dimension 3 and ghost number -1 that can be constructed with the fields ψ . The coefficients c_1 and c_2 and the functional X are taken to be independent of the scale κ . By construction, $\Gamma_R[\psi]$ generates finite Green functions since nothing in it depends on the cut-offs Λ , M_j and m_i . As regards the BRS identities, eq. (3.21) is trivially satisfied, so we only have to show that eq. (3.22) holds. That this is the case follows from eq. (3.3) and the properties $\Delta S_{\text{YM}} = 0$ and $\Delta^2 = 0$. Hence the effective action $\Gamma_R[\psi]$ with $\Gamma_R^{(1)}[\psi]$ as in eq. (3.23) defines the most general one-loop renormalized theory compatible with BRS invariance.

Note that we have defined the renormalized theory via an effective action without making reference to any bare theory. This way to proceed can be viewed as a Bogoliubov subtraction $\mathbf{T}$ on the regularized effective action, namely

$$\Gamma_R[\psi] = \lim_{\Lambda, M_j, m_i \rightarrow \infty} (1 - \mathbf{T}) \Gamma_{\Lambda M_j m_i}[\psi], \quad (3.24)$$

where

$$\mathbf{T} \Gamma_{\Lambda M_j m_i}[\psi] = \Gamma_{\Lambda m_j m_i \kappa}^{(1)\text{div}}[\psi] - c_1 S_{\text{YM}} - c_2 \Delta X[\psi]. \quad (3.25)$$

The first term on the right-hand side in eq. (3.25) removes the divergences in the regularized effective action when the regulators are sent to infinity. After subtracting this term and taking the limit $\Lambda, M_j, m_i \rightarrow \infty$, we are left with $\Gamma_{\kappa}^{(1)\text{fin}}[\psi]$. The second and third terms account for arbitrary finite BRS-invariant local renormalizations. This follows from the fact [11] that the most general solution to the equation $\Delta Y[\psi] = 0$ over the space of local integrated functionals of mass dimension 4 and ghost number zero is precisely of the form $c_1 S_{\text{YM}} + c_2 \Delta X[\psi]$. All this implies that the subtraction $\mathbf{T}$ above gives the most general BRS invariant renormalized effective action that can be defined taking as starting point the regularized effective action provided by the regularization prescription we are considering². We would like to finish this section with two comments.

Comment 1. The regularized effective action $\Gamma_{\Lambda M_j m_i}[\psi]$ from which the renormalized effective action $\Gamma_R[\psi]$ has been defined is consistent with a quantum action principle [11]. Thus the 1PI Green functions generated by $\Gamma_R[\psi]$ satisfy the renormalization group equations.

²For further details on Bogoliubov-type subtractions for regularization methods containing higher covariant derivatives see ref. [12]

Comment 2. From the point of view of Yang-Mills theory, the masses Λ , M_j and m_i are regulators on the same footing and hence there is no reason why subtraction of divergences should be done in steps. What the **T** subtraction in eqs. (3.24)-(3.25) does is precisely to subtract the divergences associated to Λ , M_j and m_i at the same time. If, say, one performs a first renormalization to subtract the divergences that occur at finite Λ when M_j and m_i are sent to infinity and then performs a second renormalization to subtract the divergences that arise when Λ goes to infinity, one finds the same of inconsistencies as in ref. [8].

4 The wrong beta function, a no-go result

In this section we compute the beta function and the anomalous dimensions of the fields. To do this, we use the renormalization group equation.

4.1 The renormalization group equations

Let us recall that if $G_R(p_e, \kappa, g, \alpha)$ denotes a renormalized 1PI Yang-Mills Green function with N_ψ external legs of type ψ , the renormalization group equation takes the form

$$\left[\kappa \frac{\partial}{\partial \kappa} + \beta(g) \frac{\partial}{\partial g} + \frac{1}{2} \sum_{\psi} \gamma_{\psi}(g) N_{\psi} + \delta(g) \frac{\partial}{\partial \alpha} \right] G_R(p_e, \kappa, g, \alpha) = 0, \quad (4.1)$$

where the coefficients $\beta(g)$, $\gamma_{\psi}(g)$ and $\delta(g)$ are power series in g ,

$$\beta(g) = \beta_1 g + \beta_2 g^2 + \dots$$

$$\gamma_{\psi}(g) = \gamma_{\psi,1} g + \gamma_{\psi,2} g^2 + \dots$$

$$\delta(g) = \delta_1 g + \delta_2 g^2 + \dots$$

The renormalization group equation is an equation in the coefficients $\beta(g)$, $\gamma_{\psi}(g)$ and $\delta(g)$ and holds for all values of the external momenta p_e^μ and the coupling constant g . We want to solve this equation up to one loop using as data the renormalized Green functions generated by the effective action $\Gamma_R[\psi]$ defined in the previous section.

So let us consider the vacuum polarization tensor generated by $\Gamma_R[\psi]$. We remind ourselves that it has the form

$$\Pi_{R\mu\nu}^{ab}(p, \kappa, g, \alpha) = - \left\{ 1 + \frac{g^2 c_V}{16\pi^2} \left[z_2(\alpha) \ln\left(\frac{p^2}{\kappa^2}\right) + \pi_0(\alpha) \right] \right\} \delta^{ab} (p^2 g_{\mu\nu} - p_\mu p_\nu) + O(g^4), \quad (4.2)$$

where for convenience we have introduced the notation

$$z_2(\alpha) = A_2 - B_2 - C_2 \quad (4.3)$$

and $\pi_0(\alpha)$ is a constant that depends on α and that already includes local finite contributions from the terms $c_1 S_{\text{YM}}$ and $c_2 \Delta X$ in eq. (3.23). Substituting $G_R(p_e, \kappa, g, \alpha)$ in eq. (4.1) by $\Pi_{R\mu\nu}^{ab}(p, \kappa, g, \alpha)$, we obtain for the left-hand side a power series in g . For the equation to hold for all g , the coefficients of this series must vanish independently. This gives to orders g , g^2 and g^3 the following three equations:

$$\begin{aligned} \gamma_{A,1} &= 0 \\ \gamma_{A,2} + \frac{c_V}{16\pi^2} \left[2\beta_1 \pi_0(\alpha) - 2z_2(\alpha) + \beta_1 \ln\left(\frac{p^2}{\kappa^2}\right) \right] &= 0 \\ \gamma_{A,3} + \frac{c_V}{16\pi^2} \left\{ 2\beta_2 \pi_0(\alpha) + \gamma_{A,1} + \delta_1 \frac{\partial \pi_0(\alpha)}{\partial \alpha} + \left[(2\beta_2 + \gamma_{A,1}) z_2(\alpha) + \delta_1 \frac{\partial z_2(\alpha)}{\partial \alpha} \right] \ln\left(\frac{p^2}{\kappa^2}\right) \right\} &= 0. \end{aligned}$$

Since the renormalization group equation holds for all values of p^μ , terms independent of p^μ and terms proportional to $\ln(p^2/\kappa^2)$ must vanish separately. This implies in particular

$$\begin{aligned} \gamma_{A,1} &= 0 & \gamma_{A,2} &= \frac{c_V}{8\pi^2} z_2(\alpha) \\ \beta_1 &= 0 & \beta_2 &= \frac{1}{2z_2(\alpha)} \frac{\partial z_2(\alpha)}{\partial \alpha} \delta_1. \end{aligned}$$

Let us next take the renormalized ghost self-energy. Our arguments in the previous section implies that it has the form

$$\Omega_R^{ab}(p, \kappa, g, \alpha) = \left\{ 1 + \frac{g^2 c_V}{16\pi^2} \left[\tilde{z}_2(\alpha) \ln\left(\frac{p^2}{\kappa^2}\right) + \omega_0(\alpha) \right] \right\} \delta^{ab} p^2 + O(g^4), \quad (4.4)$$

where $\tilde{z}_2(\alpha)$ stands for

$$\tilde{z}_2(\alpha) = \tilde{A}_2 \quad (4.5)$$

and $\omega(\alpha)$ is an α -dependent constant. Proceeding in the same way as for the vacuum polarization tensor, it is straightforward to see that the renormalization group equation for the ghost self-energy yields

$$\begin{aligned} \gamma_{c,1} + \gamma_{\bar{c},1} &= 0 & \gamma_{c,2} + \gamma_{\bar{c},2} &= \frac{c_V}{4\pi^2} \tilde{z}_2(\alpha) \\ \beta_1 &= 0 & \beta_2 &= \frac{1}{2\tilde{z}_2(\alpha)} \frac{\partial \tilde{z}_2(\alpha)}{\partial \alpha} \delta_1. \end{aligned}$$

Let us finally look at the renormalization group equation for the $\bar{c}Ac$ -vertex. If k^μ and p^μ denote the momenta of the incoming ghost and the outgoing antighost, the vertex can be written without loss of generality as

$$V_{R\mu}^{abc}(p, k, \kappa, g, \alpha) = -ig f^{abc} \left\{ \left[1 + \frac{g^2 c_V}{16\pi^2} \tilde{z}_3(\alpha) \ln\left(\frac{p^2}{\kappa^2}\right) \right] p_\mu + \frac{g^2 c_V}{16\pi^2} V_\mu^{\text{fin}}(k, p, \alpha) \right\} + O(g^5), \quad (4.6)$$

where the coefficient $\tilde{z}_3(\alpha)$ is given by

$$\tilde{z}_3(\alpha) = \tilde{A}_3 \quad (4.7)$$

and $V_\mu^{\text{fin}}(k, p, \alpha)$ contains local as well as non-local finite κ -independent radiative corrections. Using eq. (4.6) as input in the renormalization group equation, eq. (4.1), expanding in powers of g up to order two and using our previous results for β_1 and the anomalous dimensions $\gamma_{\psi,1}$ and $\gamma_{\psi,2}$, it is very to see that

$$\beta_2 = 0 \quad \beta_3 = \frac{c_V}{16\pi^2} \left[2\tilde{z}_3(\alpha) - 2\tilde{z}_2(\alpha) - z_2(\alpha) \right].$$

The first one of these equation implies in turn that

$$\delta_1 = 0.$$

Thus the renormalization group coefficients to lowest order can all be expressed in terms of $z_2(\alpha)$, $\tilde{z}_2(\alpha)$ and $\tilde{z}_3(\alpha)$. To actually compute $z_2(\alpha)$, $\tilde{z}_2(\alpha)$ and $\tilde{z}_3(\alpha)$, we must explicitly compute the limit $\Lambda, M_j, m_i \rightarrow \infty$ of the regularized vacuum polarization tensor, the regularized ghost self-energy and the regularized $\bar{c}Ac$ -vertex. We do this in the sequel.

4.2 Explicit computations

We begin with the vacuum polarization tensor $\Pi_{\mu\nu}^{ab}(p, \Lambda, M_j, m_i)$. At one loop, it receives contributions from the Feynman diagrams in Figs. 1 and 2 and in the limit $\Lambda, M_j, m_i \rightarrow \infty$ it takes the form in eqs. (3.6) and (3.7). A look at the Feynman rules in Appendix A shows that

- (i) diagrams (1e) and (2b) do not depend on any mass at all,
- (ii) diagrams (1a) and (1c) depend on Λ ,
- (iii) diagrams (1f) and (2e) depend on m_i , and
- (iv) diagrams (1b), (1d), (2a), (2c) and (2d) depend on both Λ and M_j .

This implies that the coefficient C_2 in eq. (3.7) is unambiguous, whereas the coefficients A_2 and B_2 depend on the path in (Λ, M_j) -space followed to approach $\Lambda, M_j = \infty$. Our calculations in Appendix B yield for C_2 the value

$$C_2 = -\frac{1}{3} . \quad (4.8)$$

As regards A_2 and B_2 , we consider in this paper paths of type

$$\Lambda, M_j \rightarrow \infty \quad \text{with} \quad \frac{M_j}{\Lambda} = \theta_j \quad \text{and} \quad 0 \leq \theta_j \leq \infty . \quad (4.9)$$

The path $\theta_j = \infty$ corresponds to first taking the limit $M_j \rightarrow \infty$ at finite Λ , and then taking the limit $\Lambda \rightarrow \infty$. The path $\theta_j = 0$ on the contrary corresponds to first taking the limit $\Lambda \rightarrow \infty$ at finite M_j , and then taking the limit $M_j \rightarrow \infty$. The calculations in Appendix B show that

$$\theta_j = \infty : \quad A_2 = 6 - \frac{\alpha}{2} \quad B_2 = 4 \quad (4.10)$$

$$\theta_j = 0 : \quad A_2 = -\frac{3+\alpha}{2} \quad B_2 = -\frac{21}{6} \quad (4.11)$$

$$0 < \theta_j < \infty : \quad A_2 = 2 - \frac{\alpha}{2} \quad B_2 = 0 . \quad (4.12)$$

We see that even though A_2 and B_2 are different for different θ_j , the combination $A_2 - B_2$ is the same. These results yield for $z_2(\alpha)$ in eq. (4.3)

$$z_2(\alpha) = \frac{7}{3} - \frac{\alpha}{2} . \quad (4.13)$$

It is important to note that in general the coefficients A_G , B_G and C_G are not important by themselves. What is really important are the combinations $z_G(\alpha) = A_G - B_G - C_G$, since they carry the κ -dependence of the renormalized Green functions. Note as a matter of fact that the beta function depends precisely on these combinations. In a sense, we can think of every (Λ, M_j) -path, equivalently of every set $\{A_G, B_G, C_G\}$, as defining a different regularization method for which the BRS identities (3.17)-(3.19) hold. Since we will extensively refer later on to the renormalized vacuum polarization tensor, let us put eqs. (4.2) and (4.13) together:

$$\Pi_{R\mu\nu}^{ab}(p, \kappa, g, \alpha) = - \left\{ 1 + \frac{g^2 c_V}{16\pi^2} \left[\left(\frac{7}{3} - \frac{\alpha}{2} \right) \ln \left(\frac{p^2}{\kappa^2} \right) + \pi_0(\alpha) \right] \right\} \delta^{ab} (p^2 g_{\mu\nu} - p_\mu p_\nu) . \quad (4.14)$$

Next we move on to the ghost self-energy and the $\bar{c}Ac$ -vertex. The regularized ghost self-energy at one loop $\Omega^{ab}(p, \Lambda)$ receives contributions from only the diagram in Fig. 3.

This diagram is finite by power counting and, apart from the external momentum p^μ , only depends on the mass Λ . To evaluate its limit $\Lambda \rightarrow \infty$, we use the same techniques as for the vacuum polarization tensor and obtain

$$\Omega^{(1)ab}(p, \Lambda) = \frac{g^2 c_V}{16\pi^2} \left[\frac{3-\alpha}{4} \ln\left(\frac{p^2}{\Lambda^2}\right) + \omega_0(\alpha) \right] \delta^{ab} p^2.$$

This gives for $\tilde{z}_2(\alpha)$ in eqs. (4.4)-(4.5) the value

$$\tilde{z}_2(\alpha) = \frac{3-\alpha}{4}. \quad (4.15)$$

As concerns the $\bar{c}Ac$ -vertex, at one loop it receives contributions from the two diagrams in Fig. 4. These diagrams are finite by power counting and depend on the external momenta k^μ and p^μ and the mass Λ . Using the same techniques as in Appendix B, it is not difficult to see that

$$V_\mu^{(1)abc}(k, p, \Lambda) = -ig f^{abc} \frac{g^2 c_V}{16\pi^2} \left[-\frac{\alpha}{2} \ln\left(\frac{p^2}{\Lambda^2}\right) p_\mu + \alpha V_\mu^{\text{fin}}(k, p, \alpha) \right]$$

as Λ goes to infinity. This yields

$$\tilde{z}_3 = -\frac{\alpha}{2}. \quad (4.16)$$

Substituting the values obtained for $z_2(\alpha)$, $\tilde{z}_2(\alpha)$ and $\tilde{z}_3(\alpha)$ in the expressions for the beta functions and the anomalous dimensions above, we have

$$\beta(g) = -\frac{23}{6} \frac{g^3 c_V}{16\pi^2} + O(g^4) \quad \gamma_A(g) = \left(\frac{14}{3} - \alpha\right) \frac{g^2 c_V}{16\pi^2} + O(g^3) \quad (4.17)$$

$$\gamma_c + \gamma_{\bar{c}} = (3 - \alpha) \frac{g^2 c_V}{16\pi^2} + O(g^3) \quad \delta(g) = O(g^2).$$

The results for the beta function and the anomalous dimension of the gauge field are in disagreement with the well known results

$$\beta(g) = -\frac{11}{3} \frac{g^3 c_V}{16\pi^2} + O(g^4) \quad \gamma_A(g) = \left(\frac{13}{3} - \alpha\right) \frac{g^2 c_V}{16\pi^2} + O(g^3). \quad (4.18)$$

We recall that for unitary theories the one and two-loop coefficients in the beta function and the one-loop coefficient of the anomalous dimensions are universal quantities that do not depend on the regularization method nor renormalization scheme used to compute them. In view of this, we have to conclude that the regularization prescription under consideration does not yield a consistent renormalized Yang-Mills theory, hence that it must be discarded as a candidate to regularize non-abelian gauge theories.

5 Non-physical radiative corrections from Pauli-Villars determinants

The beta function at one loop is build up with radiative corrections of type $\ln(p^2/\kappa^2)$. Hence, the fact that the result in eq. (4.17) is not correct indicates that the regularization method generates somewhere non-local logarithmic radiative corrections which should not be there. To find these corrections and their origin, we proceed as follows.

Suppose that instead of the regularization prescription of Faddeev and Slavnov we use dimensional regularization or a plain cut-off. Then the vacuum polarization tensor, the ghost self-energy and the ghost-vertex take after renormalization the form

$$\Pi'_{R\mu\nu}{}^{ab}(p, \kappa, g, \alpha) = - \left\{ 1 + \frac{g^2 c_V}{16\pi^2} \left[\left(\frac{13}{6} - \frac{\alpha}{2} \right) \ln\left(\frac{p^2}{\kappa^2}\right) + \pi_0(\alpha) \right] \right\} \delta^{ab} (p^2 g_{\mu\nu} - p_\mu p_\nu) \quad (5.1)$$

$$\Omega_R'^{ab}(p, \kappa, g, \alpha) = \left\{ 1 + \frac{g^2 c_V}{16\pi^2} \left[\frac{3-\alpha}{4} \ln\left(\frac{p^2}{\kappa^2}\right) + \omega_0(\alpha) \right] \right\} \delta^{ab} p^2$$

$$V_R' \mu^{abc}(k, p, \kappa, g, \alpha) = -ig f^{abc} \left\{ \left[1 - \frac{g^2 c_V}{16\pi^2} \frac{\alpha}{2} \ln\left(\frac{p^2}{\kappa^2}\right) \right] p_\mu + \frac{g^2 c_V}{16\pi^2} \alpha V_\mu^{\text{fin}}(k, p, \alpha) \right\}.$$

From general theorems of renormalization theory, we know that two renormalized series for a 1PI Green function that agree up to order $\hbar^{n-1}$ can differ at order $\hbar^n$ by at most a local term. It happens however that the renormalized series for the vacuum polarization tensor in eqs. (4.14) and (5.1) agree at tree level but differ at one loop by a non-local term. The question that arises then is whether eqs. (4.14) and (5.1) correspond to different but consistent quantum Yang-Mills theories. To answer this question, we remind that in Minkowski space the renormalized vacuum polarization tensor takes up to one loop the form

$$\Pi_{M\mu\nu}{}^{ab}(p, \kappa, g, \alpha) = - \left\{ 1 + \frac{g^2 c_V}{16\pi^2} \left[c(\alpha) \ln\left(\frac{-p^2 + i0^+}{\kappa^2}\right) + \pi_0(\alpha) \right] \right\} \delta^{ab} (p^2 g_{\mu\nu} - p_\mu p_\nu).$$

The coefficient $c(\alpha)$ is uniquely determined by the imaginary part of $\Pi_{M\mu\nu}{}^{ab}$,

$$\text{Im } \Pi_{M\mu\nu}{}^{ab}(p, \kappa, g, \alpha) = -c(\alpha) \frac{g^2 c_V}{16\pi^2} \theta(p^2) (p^2 g_{\mu\nu} - p_\mu p_\nu), \quad (5.2)$$

But the imaginary part of $\Pi_{M\mu\nu}{}^{ab}$ can be computed by means of Cutkosky rules as

$$\text{Im} \left(\Pi_{M\mu\nu}^{ab} \right) = \text{diagram 1} + \text{diagram 2} ,$$

where the cut propagators and the vertices are the cut propagators and vertices of unregularized Yang-Mills theory. The cut-diagrams on the right-hand side are very easy to calculate in the Feynman gauge. Doing so and using eq. (5.2), we get $c(1) = 5/3$. This selects the vacuum polarization tensor in eq. (5.1) and disregards that in eq. (4.14). It also shows that the vacuum polarization tensor as computed with higher covariant derivatives and Pauli-Villars fields contains unphysical logarithmic radiative corrections.

To pin-point the origin of these unphysical radiatives corrections, we consider all the diagrams that contribute to the vacuum polarization tensor in dimensional regularization and compute their limit $D \rightarrow 4$. Two types of contributions arise: contributions finite at $D = 4$, and contributions singular at $D = 4$. In the first ones, we set $D = 4$ and take the limit $\Lambda, M_j, m_i \rightarrow \infty$. In the second ones, we leave D arbitrary and keep the masses Λ , M_j and m_i finite. Proceeding in this way we obtain the results given in Appendix B. It follows from them that only diagrams (1a), (1e), (2a) and (2b) give $\ln(p^2)$ -contributions. The $\ln(p^2)$ -contribution from diagrams (1a) and (1e) can be read off from eqs. (B.4) and (B.6):

$$(1a) + (1e) \rightarrow - \frac{g^2 c_V}{16\pi^2} \delta^{ab} \left(\frac{13}{6} - \frac{\alpha}{2} \right) \ln(p^2) \left(p^2 g_{\mu\nu} - p_\mu p_\nu \right) . \quad (5.3)$$

As for diagrams (2a) and (2b), their $\ln(p^2)$ -contribution is [see eqs. (B.8), (B.16), (B.21) and (B.25)]

$$(2a) + (2b) \rightarrow - \frac{g^2 c_V}{16\pi^2} \delta^{ab} \frac{1}{6} \ln(p^2) \left(p^2 g_{\mu\nu} - p_\mu p_\nu \right) . \quad (5.4)$$

It thus becomes clear that diagrams (1a) and (1e) give by themselves the correct non-local part of the vacuum polarization tensor, whereas diagrams (2a) and (2b) are responsible for the unwanted logarithmic corrections. The problem is that without diagrams (2a) and (2b) the regularized vacuum polarization tensor is not finite nor transverse. This, together with the observation that diagrams (2a) and (2b) are generated by the determinants $\det \mathbb{A}_j^1$, leads to the conclusion that the very same Pauli-Villars determinants that regularize divergences also generate unphysical non-local logarithmic radiative corrections.

The determinant $\det \mathbb{A}_j^1$ has a mass Λ hidden in it. One would like to know if this

dependence on Λ plays any part in the generation of the unphysical logarithmic radiative corrections. To see if this is the case, we replace the action S_Λ in the definition of $\det \mathbb{A}_j^1$ with the Yang-Mills gauge-fixed action S in eq. (2.1) and re-compute the vacuum polarization tensor. The Feynman rules are now different and can be obtained from those in Appendix A by simply setting Λ to infinity. The diagrams that make the vacuum polarization tensor at one loop are the same as in Figs. 1 and 2, with the difference that now they are constructed with different Feynman rules. To be on the safe side, and since Pauli-Villars determinants may not prove sufficient to regularize all UV divergences, we introduce in addition dimensional regularization. Using eqs. (2.17) and (2.20), and after some lengthy but straightforward calculations, we obtain

$$\Pi_{\mu\nu}^{(1)ab}(p, M_j, m_i, \varepsilon, \mu) = -\frac{g^2 c_V}{16\pi^2} \delta^{ab} \Pi(p, M_j, m_i) (p^2 g_{\mu\nu} - p_\mu p_\nu) ,$$

where $\Pi(p, M_j, m_i, \varepsilon, \mu)$ is given by

$$\Pi(p, M_j, m_i, \varepsilon, \mu) = -\frac{3+\alpha}{2} L(p^2) - \frac{7}{2} \sum_{j=1}^J \alpha_j \ln\left(\frac{p^2}{M_j^2}\right) - \frac{1}{3} \sum_{i=1}^I \gamma_i \ln\left(\frac{p^2}{m_i^2}\right) + \pi_0(\alpha, M_j, m_i) .$$

Here $L(z^2)$ denotes

$$L(z^2) = \frac{1}{\varepsilon} + \ln\left(\frac{z^2}{\mu^2}\right) \quad D = 4 + 2\varepsilon \quad (5.5)$$

and $\pi_0(\alpha, M_j, m_i)$ collects local contributions which are finite at $D = 4$ and either vanish or remain finite as $M_j, m_i \rightarrow \infty$. The coefficient in front of $\ln(p^2)$ is $\frac{7}{3} - \frac{\alpha}{2}$, the same as in eqs. (4.2) and (4.13). Furthermore, although we have not given here partial results, it happens again that only diagrams (1a), (1e), (2a) and (2b) generate $\ln(p^2)$ -contributions and that the latter contributions have the same form as in eqs. (5.3) and (5.4). Hence the origin of the unphysical radiative corrections is not the Λ hidden in S_Λ in but the structure of the determinants $\det \mathbb{A}_j^1$.

Let us further argue that the non-local logarithmic contribution arising from diagrams (2a) and (2b) can not be set to zero by sending the masses Λ and M_j to infinity in a clever way. Suppose we take the diagrams in Figs. 1 and 2 and proceed as follows:

- (i) Set the masses Λ , M_j and m_i to infinity in the Feynman integrands and keep only those diagrams whose integrands do not vanish.
- (ii) Wick rotate to Minkowski's momentum space the surviving diagrams.
- (iii) Compute the imaginary part of the analytically continued diagrams by means of Cutkosky rules (for a general reference on cut diagrams see [13]).

Then, it is not difficult to see that only diagrams (1a), (1e), (2a) and (2b) develop an imaginary part and that in the Feynman gauge the latter takes the form

$$\begin{aligned}\text{Im} \left[(1a) + (1e) \right] &= -\frac{5}{3} \frac{g^2 c_V}{16\pi} \theta(p^2) \left(p^2 g_{\mu\nu} - p_\mu p_\nu \right) \\ \text{Im} \left[(2a) + (2b) \right] &= -\frac{1}{6} \frac{g^2 c_V}{16\pi} \theta(p^2) \left(p^2 g_{\mu\nu} - p_\mu p_\nu \right) .\end{aligned}$$

So diagrams (1a) and (1e) keep giving physical logarithmic radiative corrections and diagrams (2a) and (2b) keep giving unphysical logarithmic radiative corrections. The formal manipulations in steps (i)-(iii) are the standard manipulations used to compute the imaginary part of vacuum polarization tensor in QED [14] and rely on the following observation. In Minkowski space only diagrams with the topology of diagram (1a) develop an imaginary part at one loop. When the regulators are switched off, these diagrams are given by integrals of the form

$$\int \frac{d^4 q}{(2\pi)^4} \frac{P_{\mu\nu}(q, p)}{(q^2 + i0^+) [(q + p)^2 + i0^+]},$$

where $P_{\mu\nu}(q, p)$ is some polynomial in the momenta q^μ and p^μ . Rigorously speaking, this integral is a distribution which is not well defined. The point is that it can be properly defined without touching its imaginary part; this is what regularization and renormalization is all about. Therefore, the imaginary part of the integral before and after regularization and renormalization should be the same. Note, as a matter of fact, that for the (Λ, M_j) -paths considered in this paper, this simple way to proceed gives after Wick rotation to Euclidean space the same results as the explicit computations of Appendix B. It thus seems that the process in steps (i)-(iii) is correct if one is interested only in the imaginary part of the vacuum polarization tensor in Minkowski space, equivalently the non-local part in Euclidean space. Furthermore, since steps (i)-(iii) do not depend on the path in (Λ, M_j) -space followed to send Λ and M_j to infinity, one would expect the renormalized vacuum polarization tensor to have the form in eqs. (4.2) and (4.13) for all (Λ, M_j) -paths

The question that arises now is whether it is possible to amend the regularization method so as to get rid of the unphysical non-local radiative corrections. This is possible but at the cost of losing manifest BRS invariance. To see this, let us assume that we take the gauge non-invariant version $r = 0$ of the Pauli-Villars determinants, and let us take for concreteness the (Λ, M_j) -path in eq. (4.9) with $\theta_j = \infty$. Now there are only six Feynman diagrams that contribute to the vacuum polarization tensor, those in Fig. 1. Evaluating them in dimensional regularization and taking the limit $\Lambda, M_j, m_i \rightarrow \infty$ in those contributions which

are finite at $D = 4$, we obtain (see Appendix B for partial results)

$$\begin{aligned}
\Pi_{\mu\nu}^{(1)ab}(p, \Lambda, M_j, m_i) \Big|_{r=0} &= \frac{g^2 c_V}{16\pi^2} \delta^{ab} \left\{ -\frac{\pi\sqrt{3}}{4} \sum_{j=1}^J \alpha_j M_j^{2/3} \Lambda^{4/3} g_{\mu\nu} - \frac{1}{2} \sum_{i=1}^I \gamma_i \ln\left(\frac{m_i^2}{\mu^2}\right) m_i^2 g_{\mu\nu} \right. \\
&\quad - \left[\left(\frac{31}{18} + \frac{\alpha}{2}\right) \ln\left(\frac{p^2}{\Lambda^2}\right) + \frac{143}{36} \sum_{i=1}^I \alpha_j \ln\left(\frac{p^2}{M_j^2}\right) - \frac{1}{12} \sum_{i=1}^I \gamma_i \ln\left(\frac{p^2}{m_i^2}\right) + \text{const} \right] p^2 g_{\mu\nu} \\
&\quad \left. + \left[\left(\frac{17}{9} + \frac{\alpha}{2}\right) \ln\left(\frac{p^2}{\Lambda^2}\right) + \frac{35}{9} \sum_{i=1}^I \alpha_j \ln\left(\frac{p^2}{M_j^2}\right) + \frac{1}{6} \sum_{i=1}^I \gamma_i \ln\left(\frac{p^2}{m_i^2}\right) + \text{const} \right] p_\mu p_\nu \right\}.
\end{aligned} \tag{5.6}$$

Here we have imposed conditions (2.17), (2.20) and (2.21), which we recall from subsection 2.2.2 ensure regularization of divergences. The right-hand side in eq. (5.6) is not transverse, hence BRS invariance does not hold at the regularized level. To end up with a transverse renormalized vacuum polarization tensor, we proceed as follows. We further restrict the Pauli-Villars masses and parameters to satisfy

$$\sum_{j=1}^J \alpha_j M_j^{2/3} = 0 \quad \sum_{i=1}^I \gamma_i m_i^2 \ln(m_i^2) = 0. \tag{5.7}$$

By doing this we get rid of the first two terms in eq. (5.6). Next we introduce a renormalization mass scale κ and perform a suitable subtraction. The fact that the coefficient in front of $\ln(p^2)$ in both the terms $p^2 g_{\mu\nu}$ and $p_\mu p_\nu$ in eq. (5.6) is the same and equal to $\frac{13}{6} - \frac{\alpha}{2}$ —compare with eq. (5.1)—ensures that the renormalized vacuum polarization tensor will be transverse and will have the correct non-local part. If instead of (Λ, M_j) -paths with $\theta_j = \infty$ we take paths with $0 \leq \theta_j < \infty$ and use the results in Appendix B, it is very easy to see that the same conclusion holds. The only difference is that to ensure transversality, we must replace the condition on the masses M_j in eq. (5.7) with

$$\sum_{j=1}^J \alpha_j M_j^2 = 0.$$

It thus becomes clear that the terms introduced when $r = 0$ is replaced with $r = 1$ so as to have BRS invariance at the regularized level originate the unphysical non-local logarithmic corrections. Note in this regard that diagrams (2a) and (2b), the diagrams that originated the unphysical contributions, do not exist at all for $r = 0$. The problem with taking $r = 0$ is that manifest BRS is lost, thus becoming uninteresting.

6 Conclusion

In this paper we have addressed the question of whether higher covariant derivative Pauli-Villars regularization as proposed in refs. [2] and [3] leads to a consistent renormalized Yang-Mills theory. The regularization prescription is described by the generating functional $Z_1[J, \chi, \eta, \bar{\eta}]$ in eq. (2.12) and combines a higher covariant derivative term in the action with a set of Pauli-Villars determinants $\det A_j^1$ and $\det C_i^1$. The main results of our investigation can be summarized in the following three points:

(1) There are certain subtleties regarding the regularization mechanism that have gone unnoticed in the previous literature and that reveal that, even after imposing suitable Pauli-Villars conditions, the regularized Green functions are not finite in the power counting sense advertized in ref. [3]. This has been explicitly realized in subsection 2.2.2, where it is shown that checking that the regularized vacuum polarization tensor is finite for finite values of the regulators requires introducing a further regulator $\mathcal{R}$. This gives lie to the claim [2] [3] that the functional $Z_1[J, \chi, \eta, \bar{\eta}]$ constitutes by itself a regularized expression for the Yang-Mills path integral. Yet, since the non-local part of the regularized Green functions is the same for all admissible regulators $\mathcal{R}$, one can think of using the functional $Z_1[J, \chi, \eta, \bar{\eta}]$ to understand physical properties like *e.g.* unitarity.

(2) The regularization prescription does lead to a renormalized theory consistent with gauge invariance. This settles down some controversy in the previous literature [8] [9].

(3) The Pauli-Villars determinants on which the regularization method is based generate unphysical non-local logarithmic radiative corrections. These corrections, being non-local, survive renormalization and give an inconsistent renormalized Yang-Mills theory. In particular, the beta function of the resulting renormalized theory has a one-loop coefficient equal to $-23/6$, in contradiction with general results from renormalization theory that state that it should be $-11/3$. We expect these logarithmic unphysical corrections to break unitarity in Minkowski spacetime when the theory is coupled to matter. In fact, if unitarity is formulated in the sense of van Dam and Veltman [15], our discussion in section 5 already shows that this is the case. We have also seen that it is possible to modify the Pauli-Villars determinants so as to get rid of the unwanted radiative corrections, but this is at the expense of losing gauge invariance at the regularized level.

In view of all this we have to conclude that higher covariant derivative Pauli-Villars regulators do not provide an admissible gauge invariant regularization method.

Acknowledgements

The authors are grateful to G. 't Hooft, J.C. Taylor and M. Veltman for discussions. FRR was supported by FOM, The Netherlands. Partial support from CICyT, Spain is also acknowledged.

Appendix A: Feynman rules

Here we list the Feynman rules corresponding to the functional $Z_r[J, \chi, \eta, \bar{\eta}]$ in eq. (2.12). The propagators are given by

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{wavy line} \text{---} \bullet \\ A_\mu^a \quad A_\nu^b \end{array} = \frac{\delta^{ab} \Lambda^4}{p^4 (p^4 + \Lambda^4)} \left[p^2 g_{\mu\nu} - p_\mu p_\nu \left(1 - \frac{\alpha \Lambda^4}{p^4 + \Lambda^4} \right) \right]$$

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{wavy line} \text{---} \bullet \\ b^a \quad A_\mu^b \end{array} = -i \delta^{ab} \frac{p_\mu}{p^2}$$

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{dashed line} \text{---} \bullet \\ \bar{c}^a \quad c^b \end{array} = -\frac{\delta^{ab}}{p^2}$$

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{coiled line} \text{---} \bullet \\ A_j^a \quad A_j^b \end{array} = \frac{\delta^{ab} \Lambda^4}{p^2 (p^6 + p^2 \Lambda^4 + M_j^2 \Lambda^4)} (p^2 g_{\mu\nu} - p_\mu p_\nu)$$

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{coiled line} \text{---} \bullet \\ b_j^a \quad A_j^a \end{array} = -i \delta^{ab} \frac{p_\mu}{p^2}$$

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{double line} \text{---} \bullet \\ b_j^a \quad b_j^b \end{array} = -\delta^{ab} \frac{M_j^2}{p^2}$$

$$\begin{array}{c} p \\ \rightarrow \\ \bullet \text{---} \text{dashed line} \text{---} \bullet \\ \bar{c}_i^a \quad c_i^b \end{array} = -\frac{\delta^{ab}}{p^2 + m_i^2},$$

where p^4 stands for $(p^2)^2$. Note that the auxiliary field b_j^a propagates with propagator proportional to M_j^2 , whereas the field b^a does not propagate at all. As for the vertices, we group them according to their order in the coupling constant g . Vertices of order g are given by

$$\begin{aligned}
& \begin{array}{c} A_{\mu_1}^{a_1} \\ \downarrow p_1 \\ \bullet \\ \nearrow p_2 \quad \nwarrow p_3 \\ A_{\mu_2}^{a_2} \quad A_{\mu_3}^{a_3} \end{array} = \begin{array}{c} A_{\mu_1}^{a_1} \\ \downarrow \\ \bullet \\ \nearrow A_{j\mu_2}^{a_2} \quad \nwarrow A_{j\mu_3}^{a_3} \end{array} = \frac{ig}{\Lambda^4} \mathcal{S}_3 \left\{ f^{a_1 a_2 a_3} \left[\Lambda^4 p_{1\mu_2} g_{\mu_3 \mu_1} - p_1^4 p_{1\mu_2} g_{\mu_3 \mu_1} \right. \right. \\
& \quad \left. \left. + p_1^2 (p_3 - p_1)_{\mu_2} (p_{1\mu_3} p_{3\mu_1} - p_1 p_3 g_{\mu_3 \mu_1}) \right] \right\} \\
& \begin{array}{c} A_\mu^a \\ \downarrow \\ \bullet \\ \nearrow b_j^b \quad \nwarrow A_{j\nu}^c \end{array} = r g f^{abc} g_{\mu\nu} , \\
& \begin{array}{c} A_\mu^a \\ \downarrow \\ \bullet \\ \nearrow \bar{c}^b \quad \nwarrow c^c \end{array} = -i g f^{abc} p_\mu \\
& \begin{array}{c} A_\mu^a \\ \downarrow \\ \bullet \\ \nearrow \bar{c}_i^b \quad \nwarrow c_i^c \end{array} = -i g f^{abc} (r p_2 + p_3)_\mu
\end{aligned}$$

where $\mathcal{S}_3$ is the symmetrization operator with respect to the indices 1, 2 and 3, and r is the parameter distinguishing gauge non-invariant ($r = 0$) from gauge invariant ($r = 1$) Pauli-Villars determinants. Vertices of order g^2 have the form

The diagram shows a 4-point interaction with external momenta p_1, p_2, p_3, p_4 and external fields $A_{\mu_1}^{a_1}, A_{\mu_2}^{a_2}, A_{\mu_3}^{a_3}, A_{\mu_4}^{a_4}$. The first diagram is a tree-level exchange with a central vertex. The second diagram is a similar tree-level exchange with different internal lines. The third diagram is a loop-level exchange with a central vertex. The fourth diagram is a loop-level exchange with a central vertex. The fifth diagram is a loop-level exchange with a central vertex. The sixth diagram is a loop-level exchange with a central vertex. The seventh diagram is a loop-level exchange with a central vertex. The eighth diagram is a loop-level exchange with a central vertex. The ninth diagram is a loop-level exchange with a central vertex. The tenth diagram is a loop-level exchange with a central vertex. The eleventh diagram is a loop-level exchange with a central vertex. The twelfth diagram is a loop-level exchange with a central vertex. The thirteenth diagram is a loop-level exchange with a central vertex. The fourteenth diagram is a loop-level exchange with a central vertex. The fifteenth diagram is a loop-level exchange with a central vertex. The sixteenth diagram is a loop-level exchange with a central vertex. The seventeenth diagram is a loop-level exchange with a central vertex. The eighteenth diagram is a loop-level exchange with a central vertex. The nineteenth diagram is a loop-level exchange with a central vertex. The twentieth diagram is a loop-level exchange with a central vertex. The twenty-first diagram is a loop-level exchange with a central vertex. The twenty-second diagram is a loop-level exchange with a central vertex. The twenty-third diagram is a loop-level exchange with a central vertex. The twenty-fourth diagram is a loop-level exchange with a central vertex. The twenty-fifth diagram is a loop-level exchange with a central vertex. The twenty-sixth diagram is a loop-level exchange with a central vertex. The twenty-seventh diagram is a loop-level exchange with a central vertex. The twenty-eighth diagram is a loop-level exchange with a central vertex. The twenty-ninth diagram is a loop-level exchange with a central vertex. The thirtieth diagram is a loop-level exchange with a central vertex. The thirty-first diagram is a loop-level exchange with a central vertex. The thirty-second diagram is a loop-level exchange with a central vertex. The thirty-third diagram is a loop-level exchange with a central vertex. The thirty-fourth diagram is a loop-level exchange with a central vertex. The thirty-fifth diagram is a loop-level exchange with a central vertex. The thirty-sixth diagram is a loop-level exchange with a central vertex. The thirty-seventh diagram is a loop-level exchange with a central vertex. The thirty-eighth diagram is a loop-level exchange with a central vertex. The thirty-ninth diagram is a loop-level exchange with a central vertex. The fortieth diagram is a loop-level exchange with a central vertex. The forty-first diagram is a loop-level exchange with a central vertex. The forty-second diagram is a loop-level exchange with a central vertex. The forty-third diagram is a loop-level exchange with a central vertex. The forty-fourth diagram is a loop-level exchange with a central vertex. The forty-fifth diagram is a loop-level exchange with a central vertex. The forty-sixth diagram is a loop-level exchange with a central vertex. The forty-seventh diagram is a loop-level exchange with a central vertex. The forty-eighth diagram is a loop-level exchange with a central vertex. The forty-ninth diagram is a loop-level exchange with a central vertex. The fiftieth diagram is a loop-level exchange with a central vertex. The fifty-first diagram is a loop-level exchange with a central vertex. The fifty-second diagram is a loop-level exchange with a central vertex. The fifty-third diagram is a loop-level exchange with a central vertex. The fifty-fourth diagram is a loop-level exchange with a central vertex. The fifty-fifth diagram is a loop-level exchange with a central vertex. The fifty-sixth diagram is a loop-level exchange with a central vertex. The fifty-seventh diagram is a loop-level exchange with a central vertex. The fifty-eighth diagram is a loop-level exchange with a central vertex. The fifty-ninth diagram is a loop-level exchange with a central vertex. The sixtieth diagram is a loop-level exchange with a central vertex. The sixty-first diagram is a loop-level exchange with a central vertex. The sixty-second diagram is a loop-level exchange with a central vertex. The sixty-third diagram is a loop-level exchange with a central vertex. The sixty-fourth diagram is a loop-level exchange with a central vertex. The sixty-fifth diagram is a loop-level exchange with a central vertex. The sixty-sixth diagram is a loop-level exchange with a central vertex. The sixty-seventh diagram is a loop-level exchange with a central vertex. The sixty-eighth diagram is a loop-level exchange with a central vertex. The sixty-ninth diagram is a loop-level exchange with a central vertex. The seventieth diagram is a loop-level exchange with a central vertex. The seventy-first diagram is a loop-level exchange with a central vertex. The seventy-second diagram is a loop-level exchange with a central vertex. The seventy-third diagram is a loop-level exchange with a central vertex. The seventy-fourth diagram is a loop-level exchange with a central vertex. The seventy-fifth diagram is a loop-level exchange with a central vertex. The seventy-sixth diagram is a loop-level exchange with a central vertex. The seventy-seventh diagram is a loop-level exchange with a central vertex. The seventy-eighth diagram is a loop-level exchange with a central vertex. The seventy-ninth diagram is a loop-level exchange with a central vertex. The eightieth diagram is a loop-level exchange with a central vertex. The eighty-first diagram is a loop-level exchange with a central vertex. The eighty-second diagram is a loop-level exchange with a central vertex. The eighty-third diagram is a loop-level exchange with a central vertex. The eighty-fourth diagram is a loop-level exchange with a central vertex. The eighty-fifth diagram is a loop-level exchange with a central vertex. The eighty-sixth diagram is a loop-level exchange with a central vertex. The eighty-seventh diagram is a loop-level exchange with a central vertex. The eighty-eighth diagram is a loop-level exchange with a central vertex. The eighty-ninth diagram is a loop-level exchange with a central vertex. The ninetieth diagram is a loop-level exchange with a central vertex. The ninety-first diagram is a loop-level exchange with a central vertex. The ninety-second diagram is a loop-level exchange with a central vertex. The ninety-third diagram is a loop-level exchange with a central vertex. The ninety-fourth diagram is a loop-level exchange with a central vertex. The ninety-fifth diagram is a loop-level exchange with a central vertex. The ninety-sixth diagram is a loop-level exchange with a central vertex. The ninety-seventh diagram is a loop-level exchange with a central vertex. The ninety-eighth diagram is a loop-level exchange with a central vertex. The ninety-ninth diagram is a loop-level exchange with a central vertex. The hundredth diagram is a loop-level exchange with a central vertex.

$$\begin{aligned}
&= -\frac{g^2}{\Lambda^4} \mathcal{S}_4 \left\{ f^{a_1 a_2 b} f^{a_3 a_4 b} \left[\Lambda^4 g_{\mu_1 \mu_3} g_{\mu_2 \mu_4} + (p_1 + p_2)^4 g_{\mu_1 \mu_3} g_{\mu_2 \mu_4} \right. \right. \\
&\quad + 4 p_1^2 g_{\mu_2 \mu_3} (p_{4 \mu_1} p_{1 \mu_4} + p_1 \cdot p_4 g_{\mu_1 \mu_4}) - 4 p_1^2 g_{\mu_1 \mu_4} (2 p_1 + p_2)_{\mu_2} p_{1 \mu_3} \\
&\quad + 2 p_{1 \mu_1} p_{3 \mu_3} (p_2 \cdot p_4 g_{\mu_2 \mu_4} - p_{2 \mu_4} p_{4 \mu_2}) + 4 (p_1 + p_2)^2 g_{\mu_2 \mu_4} p_{4 \mu_1} (p_3 + 2 p_4)_{\mu_3} \\
&\quad \left. \left. + 8 (p_1 + p_2)_{\mu_1} p_{4 \mu_3} (p_2 \cdot p_4 g_{\mu_2 \mu_4} - p_{2 \mu_4} p_{4 \mu_2}) \right] \right\} \\
&= g^2 (f^{a_1 a_2 b} f^{a_4 a_3 b} + f^{a_1 a_3 b} f^{a_4 a_2 b}) g_{\mu \nu}
\end{aligned}$$

where $\mathcal{S}_4$ is the symmetrization operator with respect to the indices 1, 2, 3 and 4. The action S_Λ and $\det \mathbb{A}_j$ also contain vertices of higher order in g . However, these only enter two and higher-loop 1PI diagrams, so we do not need them here.

Appendix B: Computing the limit $\Lambda, M_j, m_i \rightarrow \infty$

In this Appendix we give the details of the calculations leading to the results of sections 4 and 5. We start presenting a large mass vanishing theorem which simplifies very much the computation of the large mass limit of the Feynman integrals we are interested in. Let us denote by $p_1^\mu, \dots, p_E^\mu$ some external momenta lying in a bounded subdomain of $\mathbb{R}^n$ and let $m > 0$ be a mass. Consider an L -loop integral of the form

$$I(p, m) = m^\eta \int d^n q_1 \dots d^n q_L \frac{M(q)}{\prod_i [(k_i^2)^{n_i} + m_i^{2n_i}]^{l_i}},$$

where η is an arbitrary real number and

$$\begin{aligned}
M(q) &= \text{monomial in the components of } q_1^\mu, \dots, q_L^\mu \\
k_i^\mu &= \text{linear combination of } p_1^\mu, \dots, p_E^\mu \text{ and } q_1^\mu, \dots, q_L^\mu \\
m_i &= \text{either } 0 \text{ or } m \\
n_i, l_i &= \text{positive integers.}
\end{aligned} \tag{B.1}$$

Given a subintegral J of $I(p, m)$, we call $\underline{\omega}_J$ to its infrared degree at vanishing external momenta and denote by $\underline{\omega}_{\min} = \min_J \{0, v\underline{\omega}_J\}$ the minimum of zero and the infrared degrees $\underline{\omega}_J$ of all the subintegrals J of $I(p, m)$ including $I(p, m)$ itself. Then, the following theorem holds:

m-Theorem. If the integral $I(p, m)$ is absolutely convergent at non-exceptional external momenta, and if its mass dimension d and $\underline{\omega}_{\min}$ defined above satisfy $d - \underline{\omega}_{\min} < 0$, then $I(p, m) \rightarrow 0$ as $m \rightarrow \infty$.

This is a trivial generalization of the m -theorem in ref. [16], to where we refer for the proof. Armed with this theorem, we proceed to derive the results in eqs. (4.10), (4.11) and (4.12) for the vacuum polarization tensor.

We start by recalling that the vacuum polarization tensor receives contributions at one loop from the eleven Feynman diagrams in Figs. 1 and 2. To compute the diagrams, we first replace each diagram by its dimensionally regularized version, then make our computations using dimensional regularization techniques, and finally take the limit $D \rightarrow 4$. To see that this way to proceed is legitimate and does not introduce ambiguities of any type, we recall the following property of dimensionally regularized integrals [17]: If I is an absolutely convergent integral defined in four dimensions, and if $I(D)$ denotes the corresponding dimensionally regularized integral, then

$$I = \lim_{D \rightarrow 4} I(D).$$

This property, together with the fact that the sums σ_1 , σ_2 and $\sigma_4^1 := (2a)$ in section 2 are given by sums of finite by power counting integrals, implies that σ_1 , σ_2 and σ_4^1 can be computed as explained. In the case of σ_3 , proving finiteness required an additional regulator $\mathcal{R}$. Our approach here takes this regulator to be dimensional regularization. It will of course happen that isolated diagrams will have poles at $D = 4$, but these poles will cancel after summing over diagrams and imposing conditions (2.17) and (2.20).

We first compute those diagrams which depend at most on one mass. To illustrate how to proceed, we take diagram (1a) as an example and look at $\alpha = 0$ contributions. With the

labeling of internal momenta in Fig. 1a, and dropping from the notation the delta δ^{ab} , the most general dimensionally regularized integral arising from diagram (1a) with $\alpha = 0$ will have the form

$$I_{\mu\nu}^{1a}(D; n_q, n_k, n_\Lambda) = \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{(\Lambda^{n_\Lambda} q^{n_q} k^{n_k} p^{n_p})_{\mu\nu}}{q^4 (q^4 + \Lambda^4) k^4 (k^4 + \Lambda^4)} . \quad (\text{B.2})$$

The numerator in the integrand is a monomial in Λ and in the components of q^μ , $k^\mu = q^\mu + p^\mu$ and p^μ of degrees n_Λ , n_q , n_k and n_p , respectively. Since the vacuum polarization tensor has mass dimension two, the n 's satisfy $n_\Lambda + n_q + n_k + n_p = 14$. Furthermore, since the propagator of the gauge field carries two powers of the momentum in the numerator, we have $n_q, n_k \geq 2$, thus ensuring IR convergence. If $n_q + n_k < 12$, the integral $I_{\mu\nu}^{1a}(D; n_q, n_k, n_\Lambda)$ is absolutely convergent by power counting at $D = 4$. In this case we can set $D = 4$ in eq. (B.2) and move on to computing its large- Λ limit. To do the latter, we use the m -theorem above. It is trivial to see that the theorem states in this case that

$$\left. \begin{array}{l} n_q + n_k + n_\Lambda < 12 \\ n_k + n_\Lambda < 8 \end{array} \right\} \Rightarrow \lim_{\Lambda \rightarrow \infty} I_{\mu\nu}^{1a}(4; n_q, n_k, n_\Lambda) = 0 .$$

If $n_q + n_k + n_\Lambda \geq 12$ and/or $n_k + n_\Lambda \geq 8$, we proceed as follows. Take *e.g.* the integral

$$J_{\mu\nu}^{1a}(D) = \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{q^{10} k^2 p_\mu q_\nu}{q^4 (q^4 + \Lambda^4) k^4 (k^4 + \Lambda^4)} . \quad (\text{B.3})$$

Substituting k^μ by $q^\mu + p^\mu$ and performing some trivial algebra, we write

$$J_{\mu\nu}^{1a}(D) = J'_{\mu\nu}(D) + J''_{\mu\nu}(D) ,$$

where

$$J'_{\mu\nu}(D) = \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{q^2 (q^2 + 2pq) p_\mu q_\nu}{(q^4 + \Lambda^4) [(q+p)^4 + \Lambda^4]}$$

and

$$J''_{\mu\nu}(D) = - \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{q^2 [p^2 q^2 - 2pq(p^2 + 2pq)] p_\mu q_\nu}{(q^4 + \Lambda^4) (q+p)^2 [(q+p)^4 + \Lambda^4]} .$$

The integral $J''_{\mu\nu}(D)$ is absolutely convergent by power counting at $D = 4$, so we can set $D = 4$ in it. Doing so and using the m -theorem, we have

$$\lim_{\Lambda \rightarrow \infty} J''_{\mu\nu}(4) = 0 .$$

To calculate $J'_{\mu\nu}(D)$, we use eq. (2.24) iteratively and whenever we come across an integral which is absolutely convergent by power counting at $D = 4$, we set $D = 4$ in it and use the

m -theorem. After two or three iterations, we get

$$J'_{\mu\nu}(D) = \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \left[\frac{q^4 p_\mu q_\nu}{(q^4 + \Lambda^4)^2} - \frac{2q^2(pq) p_\mu q_\nu}{(q^4 + \Lambda^4)^2} - \frac{4q^6(pq) p_\mu q_\nu}{(q^4 + \Lambda^4)^3} \right] + \dots,$$

with the dots collecting integrals which are finite at $D = 4$ and whose large- Λ limit at $D = 4$ vanishes. Using eq. (B.27) below, it is now trivial to see that

$$J'_{\mu\nu}(D) = \frac{3}{2} \frac{g^2 c_V}{16\pi^2} \left[\frac{1}{\varepsilon} + \ln\left(\frac{\Lambda^2}{4\pi\mu^2}\right) + \frac{1}{6} \right] p_\mu p_\nu$$

as $\varepsilon \rightarrow 0$, where $D = 4 + 2\varepsilon$. In what follows we will denote by v_0 all mass-independent contributions which are finite at $D = 4$. With this convention and the notation in eq. (5.5), we finally write

$$J_{\mu\nu}^{1a}(D) = \frac{3}{2} \frac{g^2 c_V}{16\pi^2} L(\Lambda^2) p_\mu p_\nu + v_0.$$

The contribution of any other integral (B.2) with $n_q + n_k + n_\Lambda \geq 12$ and/or $n_k + n_\Lambda \geq 8$ can be calculated in the same way. After some lengthy calculations and putting together all contributions³, we obtain for the complete diagram:

$$\begin{aligned} (1a) = & \frac{g^2 c_V}{16\pi^2} \left\{ \frac{3\pi}{2} \left(\frac{\alpha}{4} - 5 \right) \Lambda^2 g_{\mu\nu} - \left[\frac{11}{4} L(\Lambda^2) + \frac{1}{2} \left(\frac{25}{6} - \alpha \right) \ln\left(\frac{p^2}{\Lambda^2}\right) \right] p^2 g_{\mu\nu} \right. \\ & \left. - \left[13 L(\Lambda^2) - \left(\frac{7}{3} - \frac{\alpha}{2} \right) \ln\left(\frac{p^2}{\Lambda^2}\right) \right] p_\mu p_\nu \right\} + v_0. \end{aligned} \quad (B.4)$$

Diagrams (1c), (1e), (1f), (2b) and (2e) are computed in the same way. For them we obtain:

$$(1c) = \frac{g^2 c_V}{16\pi^2} \left\{ \frac{3\pi}{2} \left(5 - \frac{\alpha}{4} \right) \Lambda^2 g_{\mu\nu} + \frac{4}{3} L(\Lambda^2) \left(11 p^2 g_{\mu\nu} + p_\mu p_\nu \right) \right\} + v_0 \quad (B.5)$$

$$(1e) = -\frac{g^2 c_V}{16\pi^2} L(p^2) \left(\frac{1}{12} p^2 g_{\mu\nu} + \frac{1}{6} p_\mu p_\nu \right) + v_0 \quad (B.6)$$

$$(1f) = -\frac{g^2 c_V}{16\pi^2} \sum_{i=1}^I \gamma_i L(m_i^2) \left[\frac{(1+r)^2}{2} \mu_i^2 g_{\mu\nu} + \frac{(1+r)^2}{12} p^2 g_{\mu\nu} + \frac{1-4r+r^2}{6} p_\mu p_\nu \right] + v_0 \quad (B.7)$$

$$(2b) = \frac{g^2 c_V}{16\pi^2} r^2 \sum_{j=1}^J \alpha_j L(p^2) \left(\frac{1}{12} p^2 g_{\mu\nu} + \frac{1}{6} p_\mu p_\nu \right) + v_0 \quad (B.8)$$

$$(2e) = \frac{g^2 c_V}{16\pi^2} 2r \sum_{i=1}^I \gamma_i L(m_i^2) \mu_i^2 g_{\mu\nu} + v_0. \quad (B.9)$$

³The α -dependent part of the diagram is built up of integrals finite by power counting at $D = 4$ whose large- Λ limit can be computed using similar methods.

Let us next look at diagrams (1b), (1d), (2a), (2c) and (2d). They all depend on the masses Λ and M_j and their limit $\Lambda, M_j \rightarrow \infty$ is not unique but depends on the way in which we approach $\Lambda = M_j = \infty$. The three cases considered in section 4 are:

Case $\theta_j = \infty$

This corresponds to taking $M_j \rightarrow \infty$ ($\Lambda < \infty$ fixed) first and then sending $\Lambda \rightarrow \infty$. Let us consider diagram (1b). The most general integral it gives rise to has the form

$$I_{\mu\nu}^{1b}(D; n_q, n_k, n_\Lambda) = \sum_{j=1}^J \alpha_j \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{(\Lambda^{n_\Lambda} q^{n_q} k^{n_k} p^{n_p})_{\mu\nu}}{q^2 D_j(q) k^4 D_j(k)}, \quad (\text{B.10})$$

with

$$D_j(q) = q^6 + q^2 \Lambda^4 + M_j^2 \Lambda^4.$$

Again we have that $n_\Lambda + n_q + n_k + n_p = 14$ and $n_q, n_k \geq 2$. If $n_q + n_k < 12$, the integral is absolutely convergent by power counting at $D = 4$ and we have that

$$0 \leq |I_{\mu\nu}^{1b}(4; n_q, n_k, n_\Lambda)| \leq g^2 c_V \sum_{j=1}^J |\alpha_j| \int \frac{d^4 q}{(2\pi)^4} \frac{(\Lambda^{n_\Lambda} |q|^{n_q} |k|^{n_k} |p|^{n_p})_{\mu\nu}}{q^2 (q^6 + \lambda_j^6) k^4 (k^6 + \lambda_j^6)}, \quad (\text{B.11})$$

where

$$\lambda_j = M_j^{1/3} \Lambda^{2/3}.$$

The m -theorem above implies that the the right-hand side in eq. (B.11) goes to zero as M_j approaches infinity and Λ is kept fixed, thus yielding

$$n_q + n_k < 12 \quad \Rightarrow \quad \lim_{M_j \rightarrow \infty} I_{\mu\nu}^{1b}(4; n_q, n_k, n_\Lambda) = 0.$$

To illustrate how to proceed if $n_q + n_k \geq 12$, let us consider the integral [compare with eq. (B.3)]

$$J_{\mu\nu}^{1b}(D) = \sum_{j=1}^J \alpha_j \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{q^{10} k^2 p_\mu q_\nu}{q^2 D_j(q) k^2 D_j(k)}. \quad (\text{B.12})$$

Substituting k^μ by $q^\mu + p^\mu$ and using the identity

$$\frac{1}{D_j(q+p)} = \frac{1}{D_j(q)} - \frac{D_j(q+p) - D_j(q)}{D_j(q) D_j(q+p)} \quad (\text{B.13})$$

and the m -theorem, we get

$$J_{\mu\nu}^{1b}(D) = \sum_{j=1}^J \alpha_j \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{q^8 p_\mu q_\nu}{D_j^2(q)} \left[1 - \frac{6 q^4 (pq)}{D_j(q)} \right] + \dots$$

where the dots stand for integrals finite by power counting at $D = 4$ that vanish at $D = 4$ as $M_j \rightarrow \infty$ and $La < \infty$ is kept fixed. Recalling now

$$\frac{1}{D_j(q)} = \frac{1}{q^6 + \lambda_j^6} - \frac{q^2 \Lambda^4}{(q^6 + \lambda_j^6) D_j(q)}$$

and invoking the m -theorem again, we have

$$J_{\mu\nu}^{1b}(D) = \sum_{j=1}^J \alpha_j \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{q^8 p_\mu q_\nu}{(q^6 + \lambda_j^6)^2} \left[1 - \frac{6 q^4 (pq)}{(q^6 + \lambda_j^6)} \right] + \dots$$

Performing the integral with the help of eq. (B.28), we finally obtain

$$J_{\mu\nu}^{1b}(D) = \frac{3}{2} \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j L(\lambda_j^2) p_\mu p_\nu + v_0,$$

where the notation for $L(\lambda_j^2)$ is as in eq. (5.5). Any other integral (B.10) with $n_q + n_k \geq 12$ can be evaluated in a similar fashion. After doing so and summing over all contributions to diagram (1b), we finally get

$$(1b) = -\frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[4\pi\sqrt{3} \lambda_j^2 g_{\mu\nu} + L(\lambda_j^2) \left(\frac{11}{4} p^2 g_{\mu\nu} + 13 p_\mu p_\nu \right) \right] + v_0. \quad (B.14)$$

Note that, provided one imposes the Pauli-Villars condition (2.17), the pole in $L(\lambda_j^2)$ cancels with the pole from $L(\Lambda^2)$ in eq. (B.4) so as to give a finite partial sum σ_1 .

The other diagrams are evaluated in the same way. For themOB we obtain:

$$(1d) = \frac{1}{3} \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[\frac{23\pi\sqrt{3}}{2} \lambda_j^2 g_{\mu\nu} + 4 L(\lambda_j^2) (11 p^2 g_{\mu\nu} + p_\mu p_\nu) \right] + v_0 \quad (B.15)$$

$$(2a) = \frac{r^2}{3} \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[-\frac{\pi\sqrt{3}}{2} \lambda_j^2 g_{\mu\nu} + \ln\left(\frac{p^2}{\lambda_j^2}\right) \left(\frac{1}{4} p^2 g_{\mu\nu} - p_\mu p_\nu \right) \right] + v_0 \quad (B.16)$$

$$(2c) + (2d) = \frac{r}{3} \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[\pi\sqrt{3} \lambda_j^2 g_{\mu\nu} + 2 L(\lambda_j^2) \left(\frac{1}{4} p^2 g_{\mu\nu} - p_\mu p_\nu \right) \right] + v_0. \quad (B.17)$$

Taking $r = 1$ and summing eqs. (B.4)-(B.9) and (B.14)-(B.17), we obtain eq. (4.10). Note that, in agreement with our discussion in section 2, the partial sums σ_1 , σ_2 , σ_3^1 and $\sigma_4^1 := (2a)$ are finite (no poles) provided condition (2.17) and (2.20) are met. Note also

that contributions of type $(masses)^2 g_{\mu\nu}$ cancel in the vacuum polarization tensor upon summation. If we take $r = 0$ instead, we recover the result in eq. (5.6).

Case $\theta_j = 0$

This corresponds to first taking $\Lambda \rightarrow \infty$ at fixed M_j and then sending M_j to ∞ . Let us consider diagram (1b). As we already know, the most general dimensionally regularized Feynman integral arising from it has the form in eq. (B.10). If $n_q + n_k < 12$, the integral is finite by power counting at $D = 4$. In this case, the inequality

$$0 \leq \left| I_{\mu\nu}^{1b}(4; n_q, n_k, n_\Lambda) \right| \leq g^2 c_V \sum_{j=1}^J |\alpha_j| \int \frac{d^4 q}{(2\pi)^4} \frac{(\Lambda^{n_\Lambda} |q|^{n_q} |k|^{n_k} |p|^{n_p})_{\mu\nu}}{q^4 (q^4 + \Lambda^4) k^4 (k^4 + \Lambda^4)}$$

and the m -theorem yield

$$\left. \begin{array}{l} n_q + n_k + n_\Lambda < 12 \\ n_k + n_\Lambda < 8 \end{array} \right\} \Rightarrow \lim_{\Lambda \rightarrow \infty} I_{\mu\nu}^{1b}(4; n_q, n_k, n_\Lambda) = 0. \quad (\text{B.18})$$

To explain how to proceed if $n_q + n_k + n_\Lambda \geq 12$ and/or $n_k + n_\Lambda \geq 8$, we take again the integral $J_{\mu\nu}^{1b}(D)$ in eq. (B.12). Using eq. (2.16), eq. (B.13) and the m -theorem, it is very easy to see that

$$I_{\mu\nu}^{1b}(D) = \sum_{j=1}^J \alpha_j \frac{g^2 c_V}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \left[\frac{q^4 p_\mu q_\nu}{(q^4 + \Lambda^4)^2} - \frac{3 q^2 (pq) (2q^4 + \Lambda^4) p_\mu q_\nu}{(q^4 + \Lambda^4)^3} \right] + \dots,$$

with the dots collecting integrals finite by power counting at $D = 4$ that at $D = 4$ vanish as $\Lambda \rightarrow \infty$ and M_j is kept fixed. To perform the integration over q^μ , we employ eq. (B.27) below and obtain

$$I_{\mu\nu}^{1b}(D) = \frac{3}{2} \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j L(\Lambda^2) p_\mu p_\nu + v_0.$$

Proceeding in this way for all integrals (B.10) from diagram (1b) that escape the cut (B.18) and putting together all contributions, we end up with

$$\begin{aligned} (1b) = & \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left\{ \left[\frac{3}{4} \ln\left(\frac{p^2}{\Lambda^2}\right) - \frac{9}{4} \ln\left(\frac{\Lambda^2}{M_j^2}\right) - \frac{15}{2} \right] M_j^2 g_{\mu\nu} \right. \\ & \left. - \left[\frac{11}{4} L(\Lambda^2) - \frac{25}{12} \ln\left(\frac{\Lambda^2}{M_j^2}\right) \right] p^2 g_{\mu\nu} - \left[13 L(\Lambda^2) + \frac{7}{3} \ln\left(\frac{\Lambda^2}{M_j^2}\right) \right] p_\mu p_\nu \right\} + v_0. \end{aligned} \quad (\text{B.19})$$

For the other diagrams we get

$$(1d) = \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left\{ \left[\frac{9}{4} \ln\left(\frac{\Lambda^2}{M_j^2}\right) + \frac{15}{2} \right] M_j^2 g_{\mu\nu} + \frac{4}{3} L(\Lambda^2) (11 p^2 g_{\mu\nu} + p_\mu p_\nu) \right\} + v_0 \quad (B.20)$$

$$(2a) = r^2 \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[\frac{3}{4} \ln\left(\frac{p^2}{\Lambda^2}\right) M_j^2 g_{\mu\nu} + \frac{1}{3} \ln\left(\frac{p^2}{M_j^2}\right) \left(\frac{1}{4} p^2 g_{\mu\nu} - p_\mu p_\nu\right) \right] + v_0 \quad (B.21)$$

$$(2c) + (2d) = r^2 \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left\{ -\frac{3}{2} \ln\left(\frac{p^2}{\Lambda^2}\right) M_j^2 g_{\mu\nu} + \frac{1}{6} \left[L(\Lambda^2) + 8 \ln\left(\frac{\Lambda^2}{M_j^2}\right) \right] p^2 g_{\mu\nu} - \frac{2}{3} \left[L(\Lambda^2) + \frac{5}{4} \ln\left(\frac{\Lambda^2}{M_j^2}\right) \right] p_\mu p_\nu \right\} + v_0. \quad (B.22)$$

Eq. (4.10) follows from eqs. (B.4)-(B.9) and (B.19)-(B.22). Note again that the sums σ_l are finite and that the resulting vacuum polarization tensor is transverse.

Case $0 < \theta_j < \infty$

Using the same type of arguments methods as for the previous two cases, we have

$$(1b) = \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[a_1 \Lambda^2 g_{\mu\nu} - L(\Lambda^2) \left(\frac{11}{4} p^2 g_{\mu\nu} + 13 p_\mu p_\nu \right) \right] + v_0 \quad (B.23)$$

$$(1d) = \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[a_2 \Lambda^2 g_{\mu\nu} + \frac{4}{3} L(\Lambda^2) (11 p^2 g_{\mu\nu} + p_\mu p_\nu) \right] + v_0 \quad (B.24)$$

$$(2a) = r^2 \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[a_3 \Lambda^2 g_{\mu\nu} + \frac{1}{3} \ln\left(\frac{p^2}{\Lambda^2}\right) \left(\frac{1}{4} p^2 g_{\mu\nu} - p_\mu p_\nu\right) \right] + v_0 \quad (B.25)$$

$$(2c) + (2d) = r^2 \frac{g^2 c_V}{16\pi^2} \sum_{j=1}^J \alpha_j \left[a_4 \Lambda^2 g_{\mu\nu} + \frac{1}{3} L(\Lambda^2) \left(\frac{1}{2} p^2 g_{\mu\nu} - 2 p_\mu p_\nu\right) \right] + v_0. \quad (B.26)$$

Here $a_1, \dots, a_4$ are coefficients that depend on θ_j and that can be expressed in terms of the roots in the complex upper half-plane of the polynomial $P(z) = z^6 + z^2 + \theta_j^2$.

Useful integrals

We finish this appendix by giving the following two dimensionally regularized integrals that have been widely used in the computations and that are not in the tables:


$$\frac{1}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{(q^2)^\eta}{(q^4 + \lambda^4)^\gamma} = \frac{\lambda^{4+2\eta-4\gamma}}{32\pi^2} \left(\frac{\lambda^2}{4\pi\mu^2} \right)^{\frac{D}{2}-2} \frac{\Gamma\left(\frac{D}{4} + \frac{\eta}{2}\right) \Gamma\left(\gamma - \frac{D}{4} - \frac{\eta}{2}\right)}{\Gamma(\gamma) \Gamma\left(\frac{D}{2}\right)} \quad (\text{B.27})$$

$$\frac{1}{\mu^{D-4}} \int \frac{d^D q}{(2\pi)^D} \frac{(q^2)^\eta}{(q^6 + \lambda^6)^\gamma} = \frac{\lambda^{4+2\eta-6\gamma}}{48\pi^2} \left(\frac{\lambda^2}{4\pi\mu^2} \right)^{\frac{D}{2}-2} \frac{\Gamma\left(\frac{D}{6} + \frac{\eta}{3}\right) \Gamma\left(\gamma - \frac{D}{6} - \frac{\eta}{3}\right)}{\Gamma(\gamma) \Gamma\left(\frac{D}{2}\right)}. \quad (\text{B.28})$$


References

- [1] G. 't Hooft and M. Veltman, Nucl Phys. **B44** (1972) 189.
- [2] A.A. Slavnov, Theor. Math. Phys. **33** (1977) 977.
- [3] L.D. Faddeev and A.A. Slavnov, *Gauge fields, introduction to quantum theory*, second edition (Benjamin, Reading 1990).
- [4] Z. Bern, M.B. Halpern, L. Sadun and C. Taubes, Nucl. Phys. **284B** (1987) 35.
- [5] M. Asorey and F. Falceto, Nucl. Phys. **327B** (1989) 427.
- [6] A.A. Slavnov, *Symmetry preserving regularization for gauge and supergauge theories*, in *Superspace & supergravity*, edited by S.W. Hawking and M. Rocek, p. 177 (Cambridge University Press, Cambridge 1981).
- [7] B.J. Warr, Ann. Phys. **183** (1988) 1.
- [8] R. Sénéor, *Some remarks for the construction of Yang-Mills theories*, in *Renormalization of quantum field theories with non-linear field transformations*, edited by P. Breitenlohner, D. Maison and K. Sibold (Springer-Verlag, Berlin 1988).
- [9] M. Day, Nucl. Phys. **B213** (1983) 591.
- [10] REDUCE 3.4.1 and 3.5 (The Brand Corporation, Santa Monica 1991 and 1993).
- [11] O. Piguet and A. Rouet, Phys. Rep. **76** (1981) 1.


- [12] C.P. Martin and F. Ruiz Ruiz, *Higher covariant derivative regulators and non-multiplicative renormalization*, NIKHEF-H preprint.
- [13] G. 't Hooft and M. Veltman, *Diagrammar*, in *Particle interactions at very high energies*, edited by D. Speiser, F. Halzen and J. Weyers (Plenum Press, London 1974).
- [14] E.H. Lifshitz and L.P. Pitaevskii, *Relativistic quantum theory*, part 2, chapters XI and XII (Pergamon Press, New York 1973).
- [15] H. van Dam and M. Veltman, Nucl. Phys. **B22** (1970) 397.
- [16] G. Giavarini, C.P. Martin and F. Ruiz Ruiz, Nucl. Phys. **B381** (1992) 222.
- [17] J.C. Collins, *Renormalization* (Cambridge University Press, Cambridge 1987).




(a)




(b)




(c)



(d)




(e)




(f)


Figure 1: One-loop regularization of the vacuum polarization tensor: diagrams that contribute for both $r = 0, 1$. Recall that the contribution of (f) depends on the value of r .




(a)




(b)



(c)



(d)



(e)

Figure 2: One-loop regularization of the vacuum polarization tensor: diagrams that contribute only for $r = 1$.




Figure 3: *One-loop regularization of the ghost self-energy.*




Figure 4: *One-loop regularization of the ghost-ghost-vertex.*